

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSSPTA1644PNH

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 AUG 09 INSPEC enhanced with 1898-1968 archive  
NEWS 4 AUG 28 ADISCTI Reloaded and Enhanced  
NEWS 5 AUG 30 CA(SM)/CAplus(SM) Austrian patent law changes  
NEWS 6 SEP 21 CA/CAplus fields enhanced with simultaneous left and right  
truncation  
NEWS 7 SEP 25 CA(SM)/CAplus(SM) display of CA Lexicon enhanced  
NEWS 8 SEP 25 CAS REGISTRY(SM) no longer includes Concord 3D coordinates  
NEWS 9 SEP 25 CAS REGISTRY(SM) updated with amino acid codes for pyrrolysine  
NEWS 10 SEP 28 CEABA-VTB classification code fields reloaded with new  
classification scheme  
NEWS 11 OCT 19 LOGOFF HOLD duration extended to 120 minutes  
NEWS 12 OCT 19 E-mail format enhanced  
NEWS 13 OCT 23 Option to turn off MARPAT highlighting enhancements available  
NEWS 14 OCT 23 CAS Registry Number crossover limit increased to 300,000 in  
multiple databases  
NEWS 15 OCT 23 The Derwent World Patents Index suite of databases on STN  
has been enhanced and reloaded  
NEWS 16 OCT 30 CHEMLIST enhanced with new search and display field  
NEWS 17 NOV 03 JAPIO enhanced with IPC 8 features and functionality  
NEWS 18 NOV 10 CA/CAplus F-Term thesaurus enhanced  
NEWS 19 NOV 10 STN Express with Discover! free maintenance release Version  
8.01c now available  
NEWS 20 NOV 20 CAS Registry Number crossover limit increased to 300,000 in  
additional databases  
NEWS 21 NOV 20 CA/CAplus to MARPAT accession number crossover limit increased  
to 50,000  
NEWS 22 DEC 01 CAS REGISTRY updated with new ambiguity codes  
NEWS 23 DEC 11 CAS REGISTRY chemical nomenclature enhanced  
NEWS 24 DEC 14 WPIDS/WPINDEX/WPIX manual codes updated  
NEWS 25 DEC 14 GBFULL and FRFULL enhanced with IPC 8 features and  
functionality  
NEWS 26 DEC 18 CA/CAplus pre-1967 chemical substance index entries enhanced  
with preparation role  
NEWS 27 DEC 18 CA/CAplus patent kind codes updated  
NEWS 28 DEC 18 MARPAT to CA/CAplus accession number crossover limit increased  
to 50,000  
NEWS 29 DEC 18 MEDLINE updated in preparation for 2007 reload  
NEWS 30 DEC 27 CA/CAplus enhanced with more pre-1907 records  
  
NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS LOGIN Welcome Banner and News Items  
NEWS IPC8 For general information regarding STN implementation of IPC 8  
NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 10:06:38 ON 27 DEC 2006

=> file medline embase biosis scisearch caplus  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 10:06:55 ON 27 DEC 2006

FILE 'EMBASE' ENTERED AT 10:06:55 ON 27 DEC 2006  
Copyright (c) 2006 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 10:06:55 ON 27 DEC 2006  
Copyright (c) 2006 The Thomson Corporation

FILE 'SCISEARCH' ENTERED AT 10:06:55 ON 27 DEC 2006  
Copyright (c) 2006 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 10:06:55 ON 27 DEC 2006  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

=> s low vasopressin  
L1 53 LOW VASOPRESSIN

=> s l1 and parathyroid hormone  
L2 1 L1 AND PARATHYROID HORMONE

=> d l2 cbib abs

L2 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2006 ACS on STN

2001:31353 Document No. 134:114837 Agents for ameliorating low vasopressin level. Ogata, Etsuro; Onuma, Etsuro; Tsunenari, Toshiaki; Saito, Hidemi; Azuma, Yumiko (Chugai Seiyaku Kabushiki Kaisha, Japan). PCT Int. Appl. WO 2001002010 A1 20010111, 114 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (Japanese). CODEN: PIXXD2. APPLICATION: WO 2000-JP4413 20000703. PRIORITY: JP 1999-189322 19990702.

AB Agents for ameliorating low vasopressin level which contain as the active ingredient a substance capable of inhibiting the

binding of a parathyroid hormone-associated peptide to its receptor; and agents for ameliorating symptoms caused by a decrease in vasopressin level which contain as the active ingredient a substance capable of inhibiting the binding of a parathyroid hormone-associated peptide to its receptor.

```
=> s l1 and treatment
L3      12 L1 AND TREATMENT
```

```
=> dup remove l3
PROCESSING COMPLETED FOR L3
L4      4 DUP REMOVE L3 (8 DUPLICATES REMOVED)
```

```
=> d l4 1-4 cbib abs
```

```
L4  ANSWER 1 OF 4      MEDLINE on STN      DUPLICATE 1
2004255503.  PubMed ID: 15118420.  Sodium fraction excretion rate in
nocturnal enuresis correlates with nocturnal polyuria and osmolality.
Aceto Gabriella; Penza Rosa; Delvecchio Maurizio; Chiozza Maria Laura;
Cimador Marcello; Caione Paolo. (Department Biomedicina Eta Evolutiva,
University, Bari, Italy. ) The Journal of urology, (2004 Jun) Vol. 171,
No. 6 Pt 2, pp. 2567-70. Journal code: 0376374. ISSN: 0022-5347. Pub.
country: United States. Language: English.
AB  PURPOSE: We verify the sodium fraction excretion rate (FE Na) and
potassium fraction excretion (FE K) rates in monosymptomatic nocturnal
enuresis. We also correlate FE Na and FE K to urinary osmolality,
nocturnal polyuria and vasopressin in the same population. MATERIALS AND
METHODS: A total of 438 children 6 to 15 years old (mean age 9.7)
presenting with monosymptomatic nocturnal enuresis were recruited from
different centers. Inclusion criteria were 3 or greater wet nights a
week, no daytime incontinence and no treatment in the previous 2
months. Exclusion criteria were cardiopathy, endocrinopathy, psychiatric
problems and urinary tract abnormalities. Micturition chart, diurnal (8
am to 8 pm) and nocturnal (8 pm to 8 am) urine collection, including
separate diuresis volumes, (Na, K and Ca) electrolytes and osmolality were
evaluated, as well as serum electrolytes, creatinine and nocturnal (4 am)
vasopressin. Diurnal and nocturnal FE K and FE Na were calculated. ANOVA
test, chi-square test, Student's t test and Pearson correlation test were
used for statistical analysis. RESULTS: Nocturnal polyuria (diurnal to
nocturnal diuresis ratio less than 1) was found in 273 children (62.3%,
group 1 and nocturnal urine volumes were normal in 165 with enuresis
(37.7%, group 2). Nocturnal FE Na was abnormal in 179 children (40.8%),
including 118 in group 1 (43.2%) and 61 in group 2 (36.9%) (chi-square not
significant). FE Na was also increased in nocturnal versus daytime
diuresis (Student's t test  $p < 0.001$ ). In group 1 nocturnal FE Na
correlated with nocturnal diuresis (Pearson correlation  $p = 0.003$ ,  $r = +0.175$ ), while daytime FE Na and nocturnal FE Na correlated with diurnal
diuresis (Pearson correlation  $p = 0.001$ ,  $r = +0.225$  and Pearson
correlation  $p = 0.001$ ,  $r = +0.209$ , respectively). In group 2 nocturnal FE
Na did not correlate with diuresis (Pearson correlation  $p = 0.103$ ,  $r = +0.128$ ) but correlated with vasopressin values (Pearson correlation  $p = 0.042$ ,  $r = -0.205$ ). Urine osmolality was reduced in 140 children (31.9%)
and correlated with nocturnal diuresis (Pearson correlation  $p = 0.003$ ,  $r = -0.321$ ). Vasopressin was decreased in 332 children (75.8%, 62.6% in group
1 and 13.2% in group 2). No significant difference was found between
sexes and age of enuretic subgroups. CONCLUSIONS: Nocturnal FE Na
correlates with nocturnal diuresis, whereas daytime FE Na does not. FE K
in daytime and nighttime diuresis does not statistically differ in
nocturnal polyuric and nonpolyuric enuretic groups. Osmolality correlates
with nocturnal diuresis, and vasopressin at 4 am was lower in the
nocturnal polyuric group. The hypothesis of a subset of enuretic patients
```

presenting with nocturnal polyuria associated with high nocturnal natriuria and low vasopressin values has been confirmed.

L4 ANSWER 2 OF 4 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN DUPLICATE 2

1998:2331 Document No.: PREV199800002331. A prospective randomized trial of arginine vasopressin in the treatment of vasodilatory shock after left ventricular assist device placement. Argenziano, Michael; Choudhri, Asim F.; Oz, Mehmet C. [Reprint author]; Rose, Eric A.; Smith, Craig R.; Landry, Donald W. [Reprint author]. Milstein Hosp. Room 7-435, 177 Fort Washington Ave., New York, NY 10032, USA. Circulation, (Nov. 4, 1997) Vol. 96, No. 9 SUPPL., pp. II286-II290. print. CODEN: CIRCAZ. ISSN: 0009-7322. Language: English.

AB Background: Vasodilatory shock requiring catecholamine pressors occurs in some patients following cardiopulmonary bypass. Prompted by a clinical observation, we investigated the use of vasopressin as a treatment for this syndrome in a randomized, controlled trial. Methods and Results: Patients undergoing placement of a left ventricular assist device (n=23) were evaluated for postbypass vasodilatory shock requiring catecholamine pressors, and consecutive eligible subjects (n = 10) were evenly randomized to blinded intravenous vasopressin or saline placebo. Vasopressin (0.1 U/min) increased mean arterial pressure (57+/-4 to 84+/-2 mm Hg, P<.001) and systemic vascular resistance (813+/-113 to 1188+/-87 dyne-s/cm<sup>5</sup>, P<.001), with decreased norepinephrine administration. There was no significant response to saline, but in three subjects who crossed over, blinded vasopressin increased mean arterial pressure (69 + - 8 to 93+/-4 mm Hg) and systemic vascular resistance (898+/-88 to 1443+/-72 dyne-s/cm<sup>5</sup>) with decreased norepinephrine administration. Plasma vasopressin concentrations prior to randomization clustered in two groups: one (n=5) with concentrations inappropriately low for the degree of hypotension (8.4+/-2.1 pg/mL) and a second (n=3) with moderately elevated levels (33.7+/-1.6 pg/mL); vasopressin increased mean arterial pressure in the low vasopressin group from 57 + - 4 to 85 + 2 mm Hg (P<.01) and in the high vasopressin group from 68+/-8 to 86+/-4 mm Hg. Conclusions: Vasopressin is an effective pressor in vasodilatory shock after cardiopulmonary bypass. An absolute vasopressin deficiency was observed in the majority of patients, but all subjects responded to vasopressin administration.

L4 ANSWER 3 OF 4 MEDLINE on STN DUPLICATE 3  
1998045879. PubMed ID: 9386112. A prospective randomized trial of arginine vasopressin in the treatment of vasodilatory shock after left ventricular assist device placement. Argenziano M; Choudhri A F; Oz M C; Rose E A; Smith C R; Landry D W. (Department of Surgery, Columbia University College of Physicians and Surgeons, New York, NY, USA. ) Circulation, (1997 Nov 4) Vol. 96, No. 9 Suppl. pp. II-286-90. Journal code: 0147763. ISSN: 0009-7322. Pub. country: United States. Language: English.

AB BACKGROUND: Vasodilatory shock requiring catecholamine pressors occurs in some patients following cardiopulmonary bypass. Prompted by a clinical observation, we investigated the use of vasopressin as a treatment for this syndrome in a randomized, controlled trial. METHODS AND RESULTS: Patients undergoing placement of a left ventricular assist device (n=23) were evaluated for post-bypass vasodilatory shock requiring catecholamine pressors, and consecutive eligible subjects (n=10) were evenly randomized to blinded intravenous vasopressin or saline placebo. Vasopressin (0.1 U/min) increased mean arterial pressure (57+/-4 to 84+/-2 mm Hg, P<.001) and systemic vascular resistance (813+/-113 to 1188+/-87 dyne-s/cm<sup>5</sup>, P<.001), with decreased norepinephrine administration. There was no significant response to saline, but in three subjects who crossed over, blinded vasopressin increased mean arterial pressure (69+/-8 to 93+/-4 mm

Hg) and systemic vascular resistance ( $898 \pm 88$  to  $1443 \pm 72$  dyne-s/cm<sup>5</sup>) with decreased norepinephrine administration. Plasma vasopressin concentrations prior to randomization clustered in two groups: one (n=5) with concentrations inappropriately low for the degree of hypotension ( $8.4 \pm 2.1$  pg/mL) and a second (n=3) with moderately elevated levels ( $33.7 \pm 1.6$  pg/mL); vasopressin increased mean arterial pressure in the low vasopressin group from  $57 \pm 4$  to  $85 \pm 2$  mm Hg ( $P < .01$ ) and in the high vasopressin group from  $68 \pm 8$  to  $86 \pm 4$  mm Hg. CONCLUSIONS: Vasopressin is an effective pressor in vasodilatory shock after cardiopulmonary bypass. An absolute vasopressin deficiency was observed in the majority of patients, but all subjects responded to vasopressin administration.

- L4 ANSWER 4 OF 4 MEDLINE on STN DUPLICATE 4  
 81105539. PubMed ID: 7006294. Hypernatraemia, diabetes mellitus, hyperprolactinaemia, retarded growth and delayed puberty in a 14 year old girl. Effect of bromocriptine treatment. Christensen N C; Hagen C; Nielsen M D; Petersen S. Acta endocrinologica, (1981 Jan) Vol. 96, No. 1, pp. 30-5. Journal code: 0370312. ISSN: 0001-5598. Pub. country: Denmark. Language: English.
- AB Investigations in a 14 year old girl with arrested growth for 2 years, delayed pubertal development, hypernatraemia without thirst, diabetes mellitus and hyperlipaemia are reported. The hypernatraemia was accompanied by a low vasopressin concentration with an abnormal response to thirst, high plasma renin but normal plasma aldosterone concentrations. Treatment with vasopressin and increased fluid intake decreased serum sodium levels. Serum gonadotrophins were low; GH response during an insulin tolerance test was subnormal and basal serum Prl concentration was elevated. Bone age, thyroid function and adrenal function were normal. After initiation of bromocriptine treatment her growth accelerated and regular menstruations commenced. The serum gonadotrophin levels increased and showed pulsatile release. A hypothalamic disorder is suggested, but no cerebral lesion could be demonstrated.
- => s vasopressin deficiency  
 L5 567 VASOPRESSIN DEFICIENCY
- => s 15 and parathyroid hormone  
 L6 1 L5 AND PARATHYROID HORMONE
- => d 16 cbib abs
- L6 ANSWER 1 OF 1 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN  
 79259348 EMBASE Document No.: 1979259348. Dissociation between plasma, urine, and renal papillary cyclic AMP content following vasopressin and DDAVP. Bia M.J.; Dewitt S.; Forrest Jr. J.N.. Renal Sect., Dept. Int. Med., Yale Univ. Sch. Med., New Haven, Conn. 06510, United States. American Journal of Physiology - Renal Fluid and Electrolyte Physiology Vol. 6, No. 3, pp. F218-F225 1979. CODEN: AJPFDM  
 Pub. Country: United States. Language: English.
- AB The effects of in vivo physiologic doses of vasopressin and 1-deamino-8-D-arginine vasopressin (DDAVP) on the cyclic AMP content of plasma, urine, and renal papillary tissue were determined in the ADH-deficient Brattleboro rat. During clearance studies, plasma cyclic AMP concentrations and both total and nephrogenous urinary cyclic AMP excretion in vasopressin- and DDAVP-treated rats were similar to the values in time-matched controls. In contrast, in situ renal papillary cyclic AMP content was higher ( $P < 0.001$ ) in both vasopressin- ( $35.7 \pm$

3.6 pmol/mg protein) and DDAVP- (29.7 ± 2.2 pmol/mg protein) treated rats compared to controls (15.1 ± 1.3 pmol/mg protein). Endogenous stimulation of vasopressin by dehydration in normal rats increased both papillary cyclic AMP content (27.1 ± 2.7 pmol/mg protein) and urine osmolality, whereas no change in papillary cyclic AMP was observed following dehydration in Brattleboro rats (13.6 ± 0.8 pmol/mg protein) despite an increase in urine osmolality. The results demonstrate that changes in cyclic AMP following in vivo vasopressin are best demonstrated by measurement of in situ cyclic AMP content of the renal papilla, whereas total urinary cyclic AMP and nephrogenous cyclic AMP are not useful indices of tubular sensitivity to this hormone.

=> s 15 and hypercalcemia  
L7 0 L5 AND HYPERCALCEMIA

=> s 15 and treatment  
L8 117 L5 AND TREATMENT

=> s 18 and antibody?  
L9 0 L8 AND ANTIBOD?

=> s parathyroid hormone related protein  
L10 9998 PARATHYROID HORMONE RELATED PROTEIN

=> s 110 and PTHrP  
L11 6896 L10 AND PTHRP

=> s 111 and antibody?  
L12 857 L11 AND ANTIBOD?

=> s 112 and vasopressin  
L13 3 L12 AND VASOPRESSIN

=> dup remove l13  
PROCESSING COMPLETED FOR L13  
L14 3 DUP REMOVE L13 (0 DUPLICATES REMOVED)

=> d 114 1-3 cbib abs

L14 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2006 ACS on STN  
2001:31353 Document No. 134:114837 Agents for ameliorating low vasopressin level. Ogata, Etsuro; Onuma, Etsuro; Tsunenari, Toshiaki; Saito, Hidemi; Azuma, Yumiko (Chugai Seiyaku Kabushiki Kaisha, Japan). PCI Int. Appl. WO 2001002010 A1 20010111, 114 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (Japanese). CODEN: PIXXD2. APPLICATION: WO 2000-JP4413 20000703. PRIORITY: JP 1999-189322 19990702.

AB Agents for ameliorating low vasopressin level which contain as the active ingredient a substance capable of inhibiting the binding of a parathyroid hormone-associated peptide to its receptor; and agents for ameliorating symptoms caused by a decrease in vasopressin level which contain as the active ingredient a substance capable of inhibiting the binding of a parathyroid hormone-associated peptide to its receptor.

L14 ANSWER 2 OF 3 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights

reserved on STN

2000222942 EMBASE Parathyroid hormone-related protein as a potential target of therapy for cancer-associated morbidity. Ogata E.. Dr. E. Ogata, Cancer Institute Hospital, Japanese Found. for Cancer Research, 1-37 Kami-Ikebukuro, Toshima-ku, Tokyo 170, Japan. Cancer Vol. 88, No. 12 SUPPL., pp. 2909-2911 15 Jun 2000.

Refs: 4.

ISSN: 0008-543X. CODEN: CANCAR

Pub. Country: United States. Language: English. Summary Language: English. Entered STN: 20000713. Last Updated on STN: 20000713

AB BACKGROUND. Proinflammatory cytokines are involved in the genesis of cancer-associated cachexia. Parathyroid hormone-related protein (PTHrP) is the causative agent in humoral hypercalcemia of malignancy (HHM) and is frequently secreted from various kinds of solid tumors as well as from adult T-cell leukemia/lymphoma. PTHrP, like PTH, acts on PTH receptor type 1 (PTH1R). Activation of PTH1R may lead to stimulation of secretion of proinflammatory cytokines. It is expected, therefore, that PTHrP constitutes a key factor in the activation of the proinflammatory and cachectogenic cytokine network and consequently in the development of cachexia in patients with cancer. METHODS. Two groups of cancer-bearing patients of similar clinical backgrounds were enrolled. Plasma concentrations of PTHrP and cytokines were measured with immunoradiometric assay and radioimmunoassay, respectively. Cancer tissues from patients with HHM were transplanted into nude mice or nude rats. The effects of humanized antihuman PTHrP antibody were examined RESULTS. In clinical studies, Group B patients (with elevated plasma PTHrP), compared with Group A patients (with normal plasma PTHrP), tended to exhibit higher plasma levels of tumor necrosis factor (TNF)- $\alpha$  ( $P = 0.13$ ), interleukin(IL)-5 ( $P = 0.08$ ), and IL-8 ( $P = 0.08$ ), and had significantly higher levels of IL-6 ( $P = \leq 0.01$ ). The levels of TNF- $\alpha$  and IL-6 correlated with those of PTHrP. In animal studies, the antibody caused a prompt and sustained decline in serum calcium. This response was accompanied by improvements in food intake, drinking, body weight gain, and general behavior. It also ameliorated the suppression of serum ADH. When those effects were compared with those induced either by bisphosphonate or calcitonin, it turned out that not all of the beneficial effects of the antibody were directly correlated with the depression of blood calcium. CONCLUSIONS. PTHrP is a promising molecular target for the development of a novel mode of treatment for patients with cancer-associated morbidity. (C) 2000 American Cancer Society.

L14 ANSWER 3 OF 3 SCISEARCH COPYRIGHT (c) 2006 The Thomson Corporation on STN

1995:837036 The Genuine Article (R) Number: TJ130. PARATHYROID HORMONE-RELATED PROTEIN IS AN AUTOCRINE MODULATOR OF RABBIT PROXIMAL TUBULE CELL-GROWTH. GARCIAOCANA A (Reprint); DEMIGUEL F; PENARANDA C; ALBAR J P; SARASA J L; ESBRI P. FDN JIMENEZ DIAZ, UNIDAD METAB LAB, E-28040 MADRID, SPAIN; FDN JIMENEZ DIAZ, DEPT PATHOL, E-28040 MADRID, SPAIN; CSIC, CTR NAEL BIOTECNOL, IMMUNOL UNIT, MADRID, SPAIN. JOURNAL OF BONE AND MINERAL RESEARCH (DEC 1995) Vol. 10, No. 12, pp. 1875-1884. ISSN: 0884-0431. Publisher: BLACKWELL SCIENCE PUBL INC CAMBRIDGE, 238 MAIN ST, CAMBRIDGE, MA 02142. Language: English. \*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB Parathyroid hormone-related protein (PTHrP), a likely mediator for humoral hypercalcemia of malignancy, is also synthesized in various normal tissues. In the kidney, PTHrP, mainly detected in proximal and distal tubules, has been shown to stimulate proliferation of rat mesangial cells in culture. Experiments were carried out to investigate the

possible mitogenic effect of PTHrP in cultures of rabbit proximal tubule cells (PTC). Immunocytochemical analysis, using antihuman (h)PTHrP antibodies to (38-64) and (107-111) epitopes in the PTHrP molecule, showed strong cytoplasmic staining in PTC and in proximal tubule-like LLC-PK1 cells. PTC secreted immunoreactive PTHrP (54.8 +/- 7.0 fmol/10(6) cells) into the culture medium. Human PTHrP(1-141) stimulated proliferation in subconfluent cultures of these cells dose-dependently. This effect was similar to that induced by [Tyr(34)] hPTHrP(1-34) amide (hPTHrP[1-34]), hPTHrP(1-86), and bovine (b)PTH(1-34), while hPTHrP(38-64) amide, hPTHrP(107-111) amide, and hPTHrP(107-139) amide were ineffective. Addition of anti-hPTHrP neutralizing antibodies to (1-34), (38-64), and (107-111) epitopes of PTHrP decreased PTC growth. The mitogenic effect of these agonists was abolished in confluent PTC. In contrast, [Nle(8,18) Tyr(34)]bPTH(3-34) amide (PTH[3-34]) increased DNA synthesis in either subconfluent or confluent PTC. In LLC-PK1 cells, which also secreted PTHrP and are devoid of PTH receptors, none of these peptides affected proliferation. Forskolin (10 mu M) or H-8 (2 mu M), a protein kinase A inhibitor, did not affect basal or hPTHrP(1-34)-stimulated DNA synthesis, respectively, in subconfluent PTC. On the other hand, 10 nM staurosporine and 100 nM calphostin C, protein kinase C (PKC) inhibitors, blunted the effects of hPTHrP(1-34) or bPTH(3-34) on DNA synthesis in these cells. These studies suggest that PTHrP may function as an autocrine factor in the regulation of proximal tubule cell growth by a PKC-mediated mechanism.

```
=> s l12 and FERM BP-5631
L15      0 L12 AND FERM BP-5631
```

```
=> s FERM BP-5631
L16      0 FERM BP-5631
```

```
=> s hybridoma
L17      73304 HYBRIDOMA
```

```
=> s l17 and "FERM BP-5631"
L18      0 L17 AND "FERM BP-5631"
```

```
=> s hyperosmolarity
L19      4749 HYPEROSMOLARITY
```

```
=> s l19 and vasopressin
L20      186 L19 AND VASOPRESSIN
```

```
=> s l20 and treatment
L21      35 L20 AND TREATMENT
```

```
=> dup remove l21
PROCESSING COMPLETED FOR L21
L22      21 DUP REMOVE L21 (14 DUPLICATES REMOVED)
```

```
=> d l22 1-21 cbib abs
```

```
L22 ANSWER 1 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN
```

```
2006217915 EMBASE Plasma arginine-vasopressin following experimental stroke: Effect of osmotherapy. Chang Y.; Chen T.-Y.; Chen C.-H.; Crain B.J.; Toung T.J.K.; Bhardwaj A. A. Bhardwaj, Department of Neurology, L-226, Oregon Health and Science University, 3185 SW Sam Jackson Park Rd., Portland, OR 97239-3098, United States. bhardwaj@ohsu.edu. Journal of Applied Physiology Vol. 100, No. 5, pp.
```



1445-1451 2006.

Refs: 49.

ISSN: 8750-7587. E-ISSN: 1522-1601. CODEN: JAPHEV

Pub. Country: United States. Language: English. Summary Language: English.

Entered STN: 20060602. Last Updated on STN: 20060602

- AB Neurohumoral responses have been implicated in the pathogenesis of ischemia-evoked cerebral edema. In a well-characterized animal model of ischemic stroke, the present study was undertaken to 1) study the profile of plasma arginine-vasopressin (AVP), and 2) determine whether osmotherapy with mannitol and various concentrations of hypertonic saline (HS) solutions influence plasma AVP levels. Halothane-anesthetized adult male Wistar rats were subjected to 2 h of middle cerebral artery occlusion with the intraluminal filament technique. Plasma AVP levels (means  $\pm$  SD) were significantly elevated at 24 h ( $42 \pm 21$  pg/ml), 48 h ( $50 \pm 28$  pg/ml), and 72 h ( $110 \pm 47$  pg/ml), and returned to baseline at 96 h ( $22 \pm 15$  pg/ml) following middle cerebral artery occlusion compared with sham-operated controls ( $14 \pm 7$  pg/ml). Plasma AVP levels at 72 h were significantly attenuated with 7.5% HS ( $37 \pm 8$  pg/ml;  $360 \pm 11$  osmol/l) compared with 0.9% saline ( $73 \pm 6$ ;  $292 \pm 6$  osmol/l), 3% HS ( $66 \pm 8$  pg/ml;  $303 \pm 12$  osmol/l), or mannitol ( $74 \pm 9$  pg/ml;  $313 \pm 14$  osmol/l) treatment. HS (7.5%) significantly attenuated water content in the ipsilateral and contralateral hemispheres compared with surgical shams, 0.9% saline, 3% HS, and mannitol treatments. Peak plasma AVP levels were not associated with direct histopathological injury to the anterior hypothalamus. Attenuation of brain water content with 7.5% HS treatment coincides with attenuated serum AVP levels, and we speculate that this may represent one additional mechanism by which osmotherapy attenuates edema associated with ischemic stroke. Copyright .COPYRGIT. 2006 the American Physiological Society.

L22 ANSWER 2 OF 21 MEDLINE on STN

DUPLICATE 1

2006196896. PubMed ID: 16357093. Agonist and hypertonic saline-induced trafficking of the NK3-receptors on vasopressin neurons within the paraventricular nucleus of the hypothalamus. Haley Gwendolen E; Flynn Francis W. (Dept. of Zoology and Physiology, Univ. of Wyoming, Laramie, WY 82071, USA. ) American journal of physiology. Regulatory, integrative and comparative physiology, (2006 May) Vol. 290, No. 5, pp. R1242-50. Electronic Publication: 2005-12-15. Journal code: 100901230. ISSN: 0363-6119. Pub. country: United States. Language: English.

- AB The neurokinin 3 receptor (NK3R) is colocalized with vasopressinergic neurons within the hypothalamic paraventricular nucleus (PVN) and intraventricular injections of NK3R agonists stimulate vasopressin (VP) release. Our objectives were to test the hypotheses that intraventricular injections of the selective NK3R agonist, succinyl-[Asp6, N-Me-Phe8] substance P (senktide), activate NK3R expressed by vasopressinergic neurons within the PVN, and see whether NK3R expressed by vasopressinergic neurons in the PVN are activated by hyperosmolarity. NK3R internalization was used as a marker of receptor activation. Immunohistochemistry revealed that NK3Rs were membrane-bound on VP immunoreactive neurons in control rats. Following senktide injection, there was a significant increase in the appearance of NK3R immunoreactivity within the cytoplasm and a morphological rearrangement of the dendrites, indicating receptor internalization, which was reversible. Furthermore, pretreatment with a selective NK3R antagonist, SB-222200, blocked the senktide-induced VP release and internalization of the NK3R in the PVN. These results show that the trafficking of the NK3R is due to ligand binding the NK3R. In a subsequent experiment, rats were administered intragastric loads of 2 or 0.15 M NaCl, and NK3R immunohistochemistry was used to track activation of the receptor. In contrast to control rats, 2 M NaCl significantly increased plasma VP levels and caused the internalization of the NK3R on

VP neurons. Also, NK3R immunoreactivity was located in the nuclei of vasopressinergic neurons after senktide and 2 M NaCl treatment. These results show that hyperosmolarity stimulates the local release of an endogenous ligand in the PVN to bind to and activate NK3R on vasopressinergic neurons.

- L22 ANSWER 3 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN
- 2006295070 EMBASE Hyponatremia, hypernatremia: A physiological approach. Offenstadt G.; Das V.. G. Offenstadt, Service de Reanimation Medicale, Hopital Saint-Antoine, 184 rue du Faubourg Saint Antoine, 75012 Paris, France. georges.offenstadt@sat.aphp.fr. Minerva Anesthesiologica Vol. 72, No. 6, pp. 353-356 2006.  
Refs: 15.  
ISSN: 0375-9393. CODEN: MIANAP  
Pub. Country: Italy. Language: English. Summary Language: English; Italian.  
Entered STN: 20060711. Last Updated on STN: 20060711
- AB Natremia belongs to the toolbox of the practicing intensivist. It is an indicator of the hydration status, which is an item that must be continuously monitored in critically ill patients. Hyponatremia is not rare (1% to 2% of hospitalised patients), and hypernatremia is about 10 times less frequent while hypernatremia always indicates hypertonicity, hyponatremia is not equivalent to hypotonicity. Diagnosis depends on the history, clinical examination and basic biochemical data. It should be kept in mind that obtaining urine samples is as important as plasma samples in this respect. The first step consists in confirming that hyponatremia is hypotonic. The second step is to assess the renal response to hypotonicity. Hypotonic hyponatremia will be considered in association with hypovolemia, euvolemia or hypervolemia. The constitution of a hyperosmolar state requires an inadequate water intake. The main goal of the treatment is not to normalize numbers (they must always be checked first), but to treat symptoms. Tolerance must always be appreciated. The mathematical formulas proposed are of interest for a better understanding, but should not be followed strictly.
- L22 ANSWER 4 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN
- 2006224486 EMBASE Agonist and hypertonic saline-induced trafficking of the NK3-receptors on vasopressin neurons within the paraventricular nucleus of the hypothalamus. Haley G.E.; Flynn F.W.. F.W. Flynn, Dept. 3166, Univ. of Wyoming, 1000 E. Univ. Ave., Laramie, WY 82071, United States. flynn@uwyo.edu. American Journal of Physiology - Regulatory Integrative and Comparative Physiology Vol. 290, No. 5, pp. R1242-R1250 2006.  
Refs: 41.  
ISSN: 0363-6119. E-ISSN: 1522-1490. CODEN: AJPRDO  
Pub. Country: United States. Language: English. Summary Language: English.  
Entered STN: 20060530. Last Updated on STN: 20060530
- AB The neurokinin 3 receptor (NK3R) is colocalized with vasopressinergic neurons within the hypothalamic paraventricular nucleus (PVN) and intraventricular injections of NK3R agonists stimulate vasopressin (VP) release. Our objectives were to test the hypotheses that intraventricular injections of the selective NK3R agonist, succinyl-[Asp(6), N-Me-Phe(8)] substance P (senktide), activate NK3R expressed by vasopressinergic neurons within the PVN, and see whether NK3R expressed by vasopressinergic neurons in the PVN are activated by hyperosmolarity. NK3R internalization was used as a marker of receptor activation. Immunohistochemistry revealed that NK3Rs were membrane-bound on VP immunoreactive neurons in control rats. Following senktide injection, there was a significant increase in the appearance of NK3R immunoreactivity within the cytoplasm and a morphological

rearrangement of the dendrites, indicating receptor internalization, which was reversible. Furthermore, pretreatment with a selective NK3R antagonist, SB-222200, blocked the senktide-induced VP release and internalization of the NK3R in the PVN. These results show that the trafficking of the NK3R is due to ligand binding the NK3R. In a subsequent experiment, rats were administered intragastric loads of 2 or 0.15 M NaCl, and NK3R immunohistochemistry was used to track activation of the receptor. In contrast to control rats, 2 M NaCl significantly increased plasma VP levels and caused the internalization of the NK3R on VP neurons. Also, NK3R immunoreactivity was located in the nuclei of vasopressinergic neurons after senktide and 2 M NaCl treatment. These results show that hyperosmolarity stimulates the local release of an endogenous ligand in the PVN to bind to and activate NK3R on vasopressinergic neurons. Copyright .COPYRG. 2006 the American Physiological Society.

L22 ANSWER 5 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

2005278040 EMBASE Disorders of water imbalance. Lin M.; Liu S.J.; Lim I.T.. Dr. M. Lin, San Francisco General Hospital Emergency Services, University of California San Francisco, 1001 Potrero Avenue, San Francisco, CA 94110, United States. mlin@itsa.ucsf.edu. Emergency Medicine Clinics of North America Vol. 23, No. 3 SPEC. ISS., pp. 749-770 2005.

Refs: 92.

ISSN: 0733-8627. CODEN: EMCAD7

S 0733-8627(05)00002-7. Pub. Country: United States. Language: English. Summary Language: English.

Entered STN: 20050811. Last Updated on STN: 20050811

AB Because of the nonspecific signs and symptoms associated with disorders of water imbalance, emergency physicians must maintain a high index of suspicion for hyponatremia and hypernatremia, especially in patients at greater risk for these electrolyte disorders. Classifying patients based on their clinical volume status, serum and urine osmolality, and urine sodium concentration helps to identify the cause of the water imbalance and to tailor treatment. Specifically, important laboratory tests to order include a serum and urine sodium concentration, serum and urine osmolality, other electrolyte concentrations, and renal function tests. Choosing the appropriate type of intravenous fluid and calculating the initial fluid resuscitation rate require careful weighing of risks and benefits associated with cellular volume changes in the CNS. Correcting serum sodium concentration too slowly or too rapidly may have devastating consequences. It is essential to monitor serum sodium levels frequently, as often as every 2 to 4 hours initially, during therapy to prevent ODS in hyponatremic patients and cerebral edema in hypernatremic patients. Promising studies involving aquaretics, which are vasopressin receptor antagonists that promote free water excretion, may play a role soon. .COPYRG. 2005 Elsevier Inc. All rights reserved.

L22 ANSWER 6 OF 21 CAPLUS COPYRIGHT 2006 ACS on STN

2005:698539 Document No. 143:191391 Hyperglycemia does not increase basal hypothalamo-pituitary-adrenal activity in diabetes but it does impair the HPA response to insulin-induced hypoglycemia. Chan, Owen; Inouye, Karen; Akirav, Eitan M.; Park, Edward; Riddell, Michael C.; Matthews, Stephen G.; Vranic, Mladen (Dep. of Physiol., Unit. of Toronto, Toronto, ON, M5S 1A8, Can.). American Journal of Physiology, 289(1, Pt. 2), R235-R246 (English) 2005. CODEN: AJPHAP. ISSN: 0002-9513. Publisher: American Physiological Society.

AB Recently, we established that hypothalamo-pituitary-adrenal (HPA) and counterregulatory responses to insulin-induced hypoglycemia were impaired in uncontrolled streptozotocin (STZ)-diabetic (65 mg/kg) rats and insulin treatment restored most of these responses. In the current study, we used phloridzin to determine whether the restoration of blood glucose alone

was sufficient to normalize HPA function in diabetes. Normal, diabetic, insulin-treated, and phloridzin-treated diabetic rats were either killed after 8 days or subjected to a hypoglycemic (40 mg/dL) glucose clamp. Basal: Elevated basal ACTH and corticosterone in STZ rats were normalized with insulin but not phloridzin. Increases in hypothalamic corticotrophin-releasing hormone (CRH) and inhibitory hippocampal mineralocorticoid receptor (MR) mRNA with STZ diabetes were not restored with either insulin or phloridzin treatments. Hypoglycemia: In response to hypoglycemia, rises in plasma ACTH and corticosterone were significantly lower in diabetic rats compared with controls. Insulin and phloridzin restored both ACTH and corticosterone responses in diabetic animals. Hypothalamic CRH mRNA and pituitary pro-opiomelanocortin mRNA expression increased following 2 h of hypoglycemia in normal, insulin-treated, and phloridzin-treated diabetic rats but not in untreated diabetic rats. Arginine vasopressin mRNA was unaltered by hypoglycemia in all groups. Interestingly, hypoglycemia decreased hippocampal MR mRNA in control, insulin-, and phloridzin-treated diabetic rats but not uncontrolled diabetic rats, whereas glucocorticoid receptor mRNA was not altered by hypoglycemia. In conclusion, despite elevated basal HPA activity, HPA responses to hypoglycemia were markedly reduced in uncontrolled diabetes. We speculate that defects in the CRH response may be related to a defective MR response. It is intriguing that phloridzin did not restore basal HPA activity but it restored the HPA response to hypoglycemia, suggesting that defects in basal HPA function in diabetes are due to insulin deficiency, but impaired responsiveness to hypoglycemia appears to stem from chronic hyperglycemia.

L22 ANSWER 7 OF 21 MEDLINE on STN  
 2005069260. PubMed ID: 15698632. Protective effect of dexamethasone on osmotic-induced demyelination in rats. Sugimura Yoshihisa; Murase Takashi; Takefuji Seiko; Hayasaka Shizu; Takagishi Yoshiko; Oiso Yutaka; Murata Yoshiharu. (Department of Teratology and Genetics, Research Institute of Environmental Medicine, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan. ) Experimental neurology, (2005 Mar) Vol. 192, No. 1, pp. 178-83. Journal code: 0370712. ISSN: 0014-4886. Pub. country: United States. Language: English.

AB Central pontine myelinolysis (CPM) is a serious demyelination disease commonly associated with the rapid correction of hyponatremia. Although its pathogenesis remains unclear, the disruption of the blood-brain barrier (BBB) as a consequence of a rapid increase in serum sodium concentration is considered to play a critical role. Since glucocorticoids are known to influence BBB permeability and prevent its disruption as a result of hypertension or hyperosmolality, we investigated whether dexamethasone (DEX) could protect against osmotic demyelination in an animal model of CPM. Hyponatremia was induced in rats by liquid diet feeding and dDAVP infusion. Seven days later, the animals' hyponatremia was rapidly corrected by injecting a bolus of hypertonic saline intraperitoneally. Rats subjected to this treatment displayed serious neurological impairment and 77% died within 5 days of rapid correction of their hyponatremia; demyelination lesions were observed in various brain regions in these animals. On the other hand, rats that were treated with DEX (2 mg/kg, 0 and 6 h after hypertonic saline injection) exhibited minimal neurological impairment and all were alive after 5 days. Demyelinative lesions were rarely seen in the brains of DEX-treated rats. A marked extravasation of endogenous IgG was observed in the demyelination lesions in the brains of rats that did not receive DEX, indicating disruption of the BBB, but was not observed in DEX-treated rats. Furthermore, Evans blue injection revealed a significant reduction in staining in the brains of DEX-treated rats ( $P < 0.05$ ). These results indicate that early DEX treatment can prevent the BBB disruption that is caused by the rapid correction of hyponatremia and its associative demyelination changes, and suggest that

DEX might be effective in preventing CPM.

- L22 ANSWER 8 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN
- 2006177538 EMBASE Hyponatremia in neurological diseases in ICU. Lath R.. Dr. R. Lath, Department of Neurosurgery, Apollo Hospitals, Jubilee Hills, Hyderabad - 500 033, India. rahullath@hotmail.com. Indian Journal of Critical Care Medicine Vol. 9, No. 1, pp. 47-51 1 Jan 2005.  
Refs: 19.  
ISSN: 0972-5229. Pub. Country: India. Language: English. Summary Language: English.  
Entered STN: 20060601. Last Updated on STN: 20060601
- AB Hyponatremia is the commonest electrolyte disturbance encountered in the neurological and neurosurgical intensive care units. It can present with signs and symptoms mimicking a neurological disease and can worsen the existing neurological deficits. Hyponatremia in neurological disorders is usually of the hypoosmolar type caused either due to the Syndrome of Inappropriate Secretion of Anti Diuretic Hormone (SIADH) or Cerebral Salt Wasting Syndrome (CSWS). It is important to distinguish between these two disorders, as the treatment of the two differ to a large extent. In SIADH, the fluid intake is restricted, whereas in CSWS the treatment involves fluid and salt replacement.
- L22 ANSWER 9 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN
- 2006036347 EMBASE Diabetic hyperosmolarity: A consequence of loss of autonomy. Constans T.. T. Constans, Faculte de Medecine de Tours, Universite Francois-Rabelais, F-37032 Tours Cedex 1, France. t.constans@chu-tours.fr. Diabetes and Metabolism Vol. 31, No. SPEC. ISS. 2, pp. 5562-5566 2005.  
Refs: 22.  
ISSN: 1262-3636. CODEN: DIMEFW  
Pub. Country: France. Language: English. Summary Language: English; French.  
Entered STN: 20060209. Last Updated on STN: 20060209
- AB Diabetic hyperosmolarity is a serious acute metabolic disorder mainly occurring in the frail elderly subject presenting age-related favoring factors (reduced sensation of thirst, altered endocrine regulation), disease-related favoring factors (cognitive impairment, poor nutritional status and/or loss of autonomy), and a triggering factor, generally infection. Diabetic hyperosmolarity can occur in a previously non-diabetic patient. Intense dehydration dominates the clinical picture. The prognosis depends largely on the underlying chronic disease.
- L22 ANSWER 10 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN
- 2003497606 EMBASE Disorders of body water homeostasis. Verbalis J.G.. Prof. Dr. J.G. Verbalis, Dept. of Medicine and Physiology, Div. of Endocrinol. and Metabolism, Georgetown Univ. School of Medicine, 4000 Reservoir Road NW, Washington, DC 20007, United States. Bailliere's Best Practice and Research in Clinical Endocrinology and Metabolism Vol. 17, No. 4, pp. 471-503 2003.  
Refs: 65.  
ISSN: 1521-690X. CODEN: BBPMFY  
Pub. Country: United Kingdom. Language: English. Summary Language: English.  
Entered STN: 20031229. Last Updated on STN: 20031229
- AB Disorders of body fluids are among the most commonly encountered problems in the practice of clinical medicine. This is in large part because many different disease states can potentially disrupt the finely balanced mechanisms that control the intake and output of water and solute. It

therefore behaves clinicians treating such patients to have a good understanding of the pathophysiology, the differential diagnosis and the management of these disorders. Because body water is the primary determinant of the osmolality of the extracellular fluid, disorders of body water homeostasis can be divided into hypo-osmolar disorders, in which there is an excess of body water relative to body solute, and hyperosmolar disorders, in which there is a deficiency of body water relative to body solute. The classical hyperosmolar disorder is diabetes insipidus (DI), and the classical hypo-osmolar disorder is the syndrome of inappropriate antidiuretic hormone secretion (SIADH). This chapter first reviews the regulatory mechanisms underlying water and sodium metabolism, the two major determinants of body fluid homeostasis. The major disorders of water metabolism causing hyperosmolality and hypo-osmolality, DI and SIADH, are then discussed in detail, including the pathogenesis, differential diagnosis and treatment of these disorders.

L22 ANSWER 11 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

2003120378 EMBASE Management of disorders of water metabolism in patients with pituitary tumors. Verbalis J.G.. Dr. J.G. Verbalis, Div. of Endocrinology/Metabolism, Georgetown Univ. Sch. of Medicine, 4000 Reservoir Road NW, Washington, DC 20007, United States. verbalis@georgetown.edu. Pituitary Vol. 5, No. 2, pp. 119-132 2002. Refs: 37.

ISSN: 1386-341X. CODEN: PITUF

Pub. Country: United States. Language: English. Summary Language: English. Entered STN: 20030403. Last Updated on STN: 20030403

AB Disorders of body fluids, notably central diabetes insipidus (CDI) and the syndrome of inappropriate antidiuretic hormone secretion (SIADH), are relatively uncommon as a presenting symptom of sellar and suprasellar masses, but quite common following surgical resection of such lesions. It therefore behooves clinicians treating such patients to have a good understanding of the pathophysiology, the differential diagnosis and the management of these disorders. This review discusses some general issues concerning the pathogenesis, differential diagnosis, clinical manifestations and therapy of hyperosmolar and hyposmolar syndromes, including CDI and SIADH, and then more specifically addresses the evaluation and treatment of pre- and postoperative disorders of water metabolism in patients with pituitary adenomas.

L22 ANSWER 12 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

2001343207 EMBASE Effect of chronic treatment with haloperidol on vasopressin release and behavioral changes by osmotic stimulation of the supraoptic nucleus. Hirayama T.; Kita T.; Ogawa Y.; Ohsawa H.; Yamashita M.; Nakashima T.; Kishimoto T.. T. Kishimoto, Department of Psychiatry, Nara Medical University, 840 Shijocho, Kashihara, Nara 634-8522, Japan. toshik@naramed-u.ac.jp. Life Sciences Vol. 69, No. 18, pp. 2147-2156 21 Sep 2001. Refs: 25.

ISSN: 0024-3205. CODEN: LIFSAK

S 0024-3205(01)01295-4. Pub. Country: United States. Language: English. Summary Language: English.

Entered STN: 20011018. Last Updated on STN: 20011018

AB Chronic treatment with dopamine D(2) blockers in schizophrenic patients has been proposed as one of the causes of polydipsia and water intoxication, but this conclusion is still controversial. To investigate the relationship between dopamine D(2) blockers and these syndromes, we designed a behavioral and neurochemical study using hyperosmotic stimulation in the supraoptic nucleus (SON) by microdialysis after chronic treatment with haloperidol in rats. Animals were injected with haloperidol decanoate (20 mg/kg, i.m.) or sesame oil at 2-week intervals

for 8 successive weeks. During the 7th week, water-intake was increased 30-60 min after the hyperosmotic stimulation in both groups, but more so in haloperidol-treated animals compared to that in the control group. Moreover, arginine vasopressin (AVP) was released by the hyperosmotic stimulation in SON, but was not significantly different between groups. In addition, striatal dopamine levels 3-4 days after the microdialysis study showed a significant decrease in the haloperidol-treated animals. These results suggest that chronic treatment with haloperidol enhances water-intake produced by hyperosmotic stimulation in the SON but does not increase AVP levels in dialysates following hyperosmotic stimulation. Thus, these symptoms may be mediated by dopaminergic systems in brain. .COPYRIGHT. 2001 Elsevier Science Inc. All rights reserved.

- L22 ANSWER 13 OF 21 MEDLINE on STN DUPLICATE 2  
 1999255949. PubMed ID: 10322639. Developmental expression of urine concentration-associated genes and their altered expression in murine infantile-type polycystic kidney disease. Gattone V H 2nd; Maser R L; Tian C; Rosenberg J M; Branden M G. (Department of Anatomy and Cell Biology, University of Kansas Medical Center, Kansas City 66160-7400, USA.. vgartone@kumc.edu) . Developmental genetics, (1999) Vol. 24, No. 3-4, pp. 309-18. Journal code: 7909963. ISSN: 0192-253X. Pub. country: United States. Language: English.
- AB Currently, there is little understanding of what factors regulate the development of urine concentrating capability in normal or polycystic kidney. The present study examined the developmental expression of genes associated with urine concentration in developing mice, including C57BL/6J-cpk/cpk mice with autosomal recessive-infantile (AR) polycystic kidney disease (PKD). Concentration of urine requires: 1) medullary collecting ducts (CD) located within a hypertonic interstitium, 2) CD cell expression of functional arginine vasopressin V2 receptors (AVP-V2R), and 3) the presence of appropriate CD water channels (aquaporins, AQP 2 and 3). An increase in urine osmolality, normally seen between 1 and 3 weeks of age, was absent in cpk cystic mice. Aldose reductase mRNA expression (a gene upregulated by medullary hyperosmolality) increased in normal mice, but remained low in the cystic kidney, suggesting the absence of a hypertonic medullary interstitium. AVP-V2R, AQP2, and AQP3 mRNA expression normally increase between 7 and 14 days. However, all were dramatically overexpressed even at 7 days of age in the cpk kidney in vivo, but decreased in vitro. Activation of the AVP-V2 receptor stimulates the production of cAMP, a substance known to promote cyst enlargement. To determine if CD cAMP, generated from increased AVP-V2Rs, was accelerating the PKD, cystic mice and their normal littermates were treated with OPC31260, a relatively specific AVP-V2R antagonist. OPC31260 treatment of cystic mice led to an amelioration of the cystic enlargement and azotemia. Treatment also decreased renal AQP2 mRNA but increased AVP-V2R and AQP3 mRNA expression in vivo. AVP upregulates the expression of AVP-V2R, AQP2, and AQP3 mRNAs in vitro. Renal EGF, known to inhibit AVP-V2R activity, downregulates AVP-V2R mRNA in vitro. Brief in vivo EGF treatment, known to decrease PKD in cpk mice, led to increased expression of AVP-V2R, AQP2, and AQP3 mRNAs at 2 weeks in both normal and cystic mice but no change was evident at 3 weeks of age. In conclusion, the development of urinary concentration ability correlates with the development of an increased medullary osmotic gradient which is diminished in murine ARPKD. However, CD genes associated with this process are overexpressed in vivo but underexpressed in vitro in the cystic kidney. The overexpression and/or overactivity of the AVP-V2R appears to contribute to the progression of PKD since an AVP-V2R antagonist inhibits cystic renal enlargement in the cpk mouse.

reserved on STN

DUPLICATE 3

1998188767 EMBASE Upregulation of hypothalamic nitric oxide synthase gene expression in streptozotocin-induced diabetic rats. Serino R.; Ueta Y.; Tokunaga M.; Hara Y.; Nomura M.; Kabashima N.; Shibuya I.; Hattori Y.; Yamashita H.. Dr. H. Yamashita, Department of Physiology, School of Medicine, Univ. of Occupational/Envntl. Health, 1-1 Iseigaoka, Yahatanishi-ku, Kitakyushu 807-8555, Japan. Diabetologia Vol. 41, No. 6, pp. 640-648 1998.

Refs: 31.

ISSN: 0012-186X. CODEN: DBTGJAJ

Pub. Country: Germany. Language: English. Summary Language: English.

Entered STN: 19980709. Last Updated on STN: 19980709

AB Plasma arginine vasopressin (AVP) is known to be elevated in patients with uncontrolled insulin-dependent diabetes mellitus who have plasma hyperosmolality with hyperglycaemia. Although osmotic stimuli cause an increase in nitric oxide synthase (NOS) activity as well as synthesis of AVP and oxytocin in the paraventricular (PVN) and supraoptic nuclei (SON), it is not known whether NOS activity in the hypothalamus changes in the diabetic patients who have plasma hyperosmolality with hyperglycaemia caused by insulin deficiency. Expression of the neuronal (n) NOS gene in the PVN and SON in streptozotocin (STZ)-induced diabetic rats was investigated by using *in situ* hybridization histochemistry and NADPH-diaphorase histochemical staining. Four weeks after intraperitoneal (i. p.) administration of STZ, male Wistar rats developed hyperglycaemia and plasma hyperosmolality. The expression of nNOS gene and NADPH-diaphorase staining in the PVN and SON remarkably increased in STZ-induced diabetic rats compared to control rats. Three weeks after administration of STZ, the diabetic rats were subcutaneously treated with insulin for 1 week, which resulted in significant suppression of the induction of nNOS, AVP and oxytocin gene expression in the PVN and SON. Furthermore, the induction of nNOS gene expression in the PVN and SON was suppressed in STZ-induced diabetic rats treated with phlorizin and diet to normalize hyperglycaemia without insulin treatment. These results suggest that upregulation of nNOS gene expression as well as AVP and oxytocin gene expression in the PVN and SON in STZ-induced diabetic rats may be associated with hyperglycaemia and plasma hyperosmolality.

L22 ANSWER 15 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

1998309509 EMBASE Hypertonic saline resuscitation. Rocha e Silva M.. Dr. M. Rocha e Silva, Instituto del Corazon, Av. Eneas de Carvalho Aguiar 44, Sao Paulo, SP, CEP 05403-000, Brazil. mrsilva@incor.usp.br. Medicina Vol. 58, No. 4, pp. 393-402 1998.

Refs: 186.

ISSN: 0025-7680. CODEN: MEDCAD

Pub. Country: Argentina. Language: English. Summary Language: English; Spanish.

Entered STN: 19981009. Last Updated on STN: 19981009

AB Treatment of severe hemorrhage offers few theoretical problems, but in practice, severe blood loss usually occurs out of hospital, often in more or less inaccessible scenarios. Controversy rages over ideal fluid, ideal volume, and minimum O<sub>2</sub> carrying capacity, but all agree that pre-hospital, isotonic resuscitation is unfeasible. The effects of highly hypertonic 7.5% NaCl (HS) was first described in 1980, when we showed that it induced immediate and long lasting hemodynamic restoration. The addition of 6% dextran-70 to (HSD) significantly enhances the duration and intensity of volume expansion, with no loss of hemodynamic effects. HS/HSD restores cardiac output, arterial pressure, base excess and oxygen availability, induce pre-capillary vasodilation, moderate hyperosmolality and hypernatremia, reversal of high glucose and lactate. It interferes with endocrine secretions when administered to animals in hemorrhagic hypotension. HS acts through transient plasma



volume expansion, positive inotropic effect on cardiac contractility, precapillary vasodilation through a direct action on vascular smooth muscle. Expansion of circulating volume is part of the mechanism, the extra volume coming from the intracellular compartment fluid, especially from endothelial and red blood cells, which facilitate microcirculatory flow. The new field of interactions of hypertonicity with the immune mechanisms may provide insight into the long lasting effects of hypertonic solutions. Randomized double blind prospective studies on the effects of HS, or HSD, used as first treatment of shock show that both are safe and free from collateral, toxic effects. These studies show an early significant rise in arterial blood pressure and a non-significant trend towards higher levels of survival. HSD administration to patients about to undergo cardiopulmonary bypass for cardiac surgery results in higher cardiac output before, and immediately following cardiopulmonary bypass, as well as zero fluid balance.

- L22 ANSWER 16 OF 21 MEDLINE on STN DUPLICATE 4  
 1998070398. PubMed ID: 9405432. Hyperosmolarity-induced gene stimulation is mediated by the negative calcium responsive element. Okazaki T; Ishikawa T; Nishimori S; Igarashi T; Hata K; Fujita T. (Endocrine Genetics and Hypertension Unit, 4th Department of Internal Medicine, University of Tokyo School of Medicine, Bunkyo-ku, Tokyo 112, Japan.. okbgenei-ky@umin.u-tokyo.ac) . The Journal of biological chemistry, (1997 Dec 19) Vol. 272, No. 51, pp. 32274-9. Journal code: 2985121R. ISSN: 0021-9258. Pub. country: United States. Language: English.
- AB The negative calcium responsive elements of the parathyroid hormone gene bind to a specific set of nuclear proteins in an extracellular calcium (Ca<sup>2+</sup>) -dependent manner. We have found that one of the negative calcium responsive elements, named oligo B, is found in the 5'-flanking region of such vasoactive genes as the vasopressin and atrial natriuretic polypeptide genes. Furthermore, the oligo B-like sequence in the former gene is conserved throughout evolution. Because expression of some of these vasoactive genes is altered by external stimuli which change cell volume, we examined whether oligo B is involved in gene regulation by hyperosmolarity. Here, we demonstrate that the binding between oligo B and its binding nuclear proteins including a redox factor 1 was reduced by hyperosmolarity generated by sodium chloride but not by urea. Such attenuated binding was reversed by dephosphorylating nuclear proteins by a potato acid phosphatase, suggesting that NaCl treatment elicited phosphorylation of these nuclear proteins to weaken their binding activity to oligo B. Furthermore, these nuclear events led to hyperosmolarity-mediated transcriptional stimulation of the genes bearing this DNA element in the cultured cells.
- L22 ANSWER 17 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN DUPLICATE 5  
 88027910 EMBASE Document No.: 1988027910. Chronic intracerebroventricular morphine and lactation in rats: Dependence and tolerance in relation to oxytocin neurones. Rayner V.C.; Robinson I.C.A.F.; Russell J.A.. Department of Physiology, University Medical School, Edinburgh, EH8 9AG, United Kingdom. Journal of Physiology Vol. 396, pp. 319-347 1988. ISSN: 0022-3751. CODEN: JPHYA7  
 Pub. Country: United Kingdom. Language: English. Summary Language: English.  
 Entered STN: 911211. Last Updated on STN: 911211
- AB 1. Acutely, opioids inhibit oxytocin secretion. To study the responses of oxytocin neurones during chronic opioid exposure, forty-five lactating rats were infused continuously from a subcutaneous osmotically driven mini-pump via a lateral cerebral ventricle with morphine sulphate solution from day 2 post-partum for 5-7 days; the infusion rate was increased 2- or 2.5-fold each 40 h from 10 µg/h initially up to 50 µg/h; controls were infused with vehicle (1 µl/h, twenty-eight rats) or were untreated

(eight rats). 2. Maternal behaviour was disrupted in 27% of the morphine-treated rats; in rats that remained maternal morphine did not affect body weight or water intake but increased rectal temperature by  $0.82 \pm 0.14^{\circ}\text{C}$  (mean  $\pm$  S.E.M.) across the first 4 days. 3. Weight gain of the litters of maternal morphine-treated rats was reduced by 32% during 7 days, predominantly in the first day of treatment when milk transfer was also reduced. Observation of pup behaviour during suckling showed decreased frequency of milk ejections on only the second day of morphine treatment. Plasma concentration of prolactin after 6 days was similar in maternal morphine-treated and control rats, but reduced by 90% in non-maternal morphine-treated rats, indicating normal control of prolactin secretion by suckling in morphine-treated rats. 4. Oxytocin and vasopressin contents, measured by radioimmunoassay, in the supraoptic and paraventricular nuclei and in the neurohypophysis were similar between fourteen maternal morphine-treated, twelve vehicle-treated and eight untreated lactating rats; thus exposure to morphine did not involve increased production and storage of oxytocin. 5. Distribution of [ $^3\text{H}$ ]morphine infused intracerebroventricularly into six virgin female rats for 6 days was measured by scintillation counting of tissue extracts. Morphine concentration in the hypothalamus and neurohypophysis was 2.7 and 12.8  $\mu\text{g/g}$ , respectively, and in blood plasma 0.75  $\mu\text{g/g}$ . Tolerance was not due to failure of morphine infusion. In addition, naloxone (5 mg/kg S.C.) provoked typical withdrawal reactions ('wet dog' shakes, defaecation, burrowing) in lactating rats infused with morphine for 5 days. 6. Pups were suckled onto seven maternal morphine-infused and five vehicle-infused rats anaesthetized with urethane for recording of intramammary and arterial blood pressures after treatment for 5 days. The incidence and pattern of milk ejections, and mammary gland sensitivity to oxytocin were similar in the two groups. Tolerance to the inhibition of suckling-induced oxytocin secretion by intracerebroventricular (I.C.V.) morphine did not extend to acute intravenous morphine (2.5 or 5 mg/kg). 7. In fourteen out of fifteen morphine-infused rats under urethane anaesthesia, intravenous naloxone HCl (5 mg/kg) quickly provoked a large, fluctuating increase in intramammary pressure lasting  $41.5 \pm 15$  min (mean  $\pm$  S.D.); this excitation of presumptive oxytocin secretion was independent of suckling and was not seen in twelve vehicle-infused rats. Carbachol I.C.V. (0.2  $\mu\text{g}$ ) produced a similar excitation of oxytocin release in both groups of rats. Naloxone did not reveal or produce stimulation of oxytocin secretion by hypotension, hypernatraemia or hyperosmolality of extracellular fluid. 8. It is concluded that chronic I.C.V. morphine infusion leads to the development of both tolerance and dependence in the mechanisms that lead to oxytocin secretion, i.e. in an identifiable peptidergic neurosecretory system. This provides a new model for the study of the processes involved in the acute and chronic actions of opioids on neurones.

L22 ANSWER 18 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

86058034 EMBASE Document No.: 1986058034. [The principal endocrine paraneoplastic syndromes accompanying lung cancer. Pathophysiologic-clinical correlations and therapeutic approaches]. LE PRINCIPALI SINDROMI PARANEOPLASTICHE ENDOCRINE NEL CANCRO POLMONARE. CORRELAZIONI FISIOPATOLOGICO-CLINICHE ED APPROCCIO TERAPEUTICO. Di Lollo F.; Cerinic M.M.. Divisione Medica, USL N. 10, Ospedale di Careggi, Istituto di Clinica Medica II dell'Universita di Firenze, Firenze, Italy. Clinica Terapeutica Vol. 115, No. 4, pp. 283-295 1985. CODEN: CLTEA4

Pub. Country: Italy. Language: Italian. Summary Language: English. Entered STN: 911210. Last Updated on STN: 911210

AB Discussion is limited to the most frequently observed paraneoplastic syndromes accompanying lung cancer, among which inappropriate ACTH

secretion is the most frequent one. From the clinical point of view certain aspects of this syndrome differ from Cushing syndrome in as much as it is characterized by lack of response of urinary 17 KS to tests of stimulation and inhibition, by high levels of circulating ACTH and failure of dexamethasone administration to influence cortisol blood level. Therapeutic intervention is aimed above all at the speedy correction of K<sup>+</sup> depletion. The paraneoplastic syndrome due to inadequate ADH secretion is characterized by fluid retention (increased circulating blood volume), hyponatremia and hypochloremia (plasma hyperosmolality), oliguria and absence of heart, kidney and adrenal disorders. Therapeutic measures consist in limiting fluid intake, administering cortisone, and adding NaCl to slow infusion only in cases with signs of cerebral impairment. Pseudohyperparathyroidism caused by the secretion of PTH-similar substances is accompanied by high urinary hydroxyproline excretion and disorders of phosphorus and calcium household: hypercalcemia, increased alkaline phosphatase, increased calciuria and phosphaturia. Hypercalcemia requires early treatment using calcitonin in order to reduce circulating Ca<sup>++</sup> (blockage of osteoclasts), corticosteroids (block of Ca<sup>++</sup> absorption in the intestine), furosemide (increased urinary Ca<sup>++</sup> excretion). In addition, fluid and salt depletion and the metabolic deviations favoring calcium ionization must be corrected.

- L22 ANSWER 19 OF 21 MEDLINE on STN DUPLICATE 6  
 82132708. PubMed ID: 7036730. Neurogenic disorders of osmoregulation. Robertson G L; Aycinena P; Zerbe R L. The American journal of medicine, (1982 Feb) Vol. 72, No. 2, pp. 339-53. Ref: 110. Journal code: 0267200. ISSN: 0002-9343. Pub. country: United States. Language: English.
- AB The osmolality of body fluids is normally maintained within a narrow range. This constancy is achieved largely via hypothalamic osmo-receptors that regulate thirst and arginine vasopressin, the antidiuretic hormone (ADH). Anything that interferes with the full expression of either osmoregulatory function exposes the patient to the hazards of abnormal increases or decreases in plasma osmolality. Hyposmolality is almost always due to a defect in water excretion. Increased intake may contribute to the problem but is rarely, if ever, a sufficient cause. Impaired water excretion can be due to a primary defect in the osmoregulation of ADH (inappropriate antidiuresis) or secondary to nonosmotic stimuli like hypovolemia or nausea. The two types differ in clinical presentation and treatment. Resetting of the ADH osmostat is commonly associated with resetting of the thirst osmostat. Hyperosmolality is almost always due to deficient water intake. Excessive excretion may contribute to the problem but is never a sufficient cause. Impaired water intake can result from a defect in either the osmoregulation of thirst of the necessary motor responses. Thirst may be deficient because of primary osmoreceptor damage as in the syndrome of adipsic hypernatremia or secondary to nonosmotic influences on the set of the system. They are distinguishable by the clinical presentation as well as the type of ADH defects with which they are associated. So-called essential hypernatremia due to primary resetting of the osmostat has been postulated, but unambiguous evidence for such an entity has not yet been reported.
- L22 ANSWER 20 OF 21 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN  
 79145974 EMBASE Document No.: 1979145974. [Hypothalamic hyperosmolality in childhood]. HYPOTHALAMISCH BEDINGTE STORUNGEN DER OSMOREGULATION IM KINDESALTER. Andler W.; Roosen K.; Reinhardt V.. Klinderklin., Univ., GHS Essen, Germany. Neurochirurgia Vol. 22, No. 2, pp. 56-68 1979.  
 CODEN: NURABV  
 Pub. Country: Germany. Language: German. Summary Language: English.

AB Hypothalamic lesions occasionally lead to excessive hypernatraemia and hyperosmolality which cannot be explained by defective ADH secretion alone. As osmoregulation is a complex system the clinical features differ widely from one patient to another. In general central dysregulation of osmolality is due to diffuse hypothalamic lesions, e.g. inflammatory infiltration by histiocytosis X or by large suprasellar tumours. We report on a ten-year-old girl suffering from a suprasellar spongioblastoma and a twelve-year-old girl, who had been operated on for a large craniopharyngioma. Polyuria and polydipsia were not present. Whereas one patient presented hypernatraemic crises and showed normal osmolality at the intervals, the other patient suffered from sustained hypernatraemia and hyperosmolality. In the first patient water loading led promptly to clinical and laboratory normalisation. In the other case water loading failed to decrease hyperosmolality but led to oedema. In the first patient hypernatraemic crises were combined with decreased serum potassium levels and elevated urinary aldosterone excretion. Therefore acute and long-term trials of spironolactone treatment were successful. Exogenous ADH-derivates failed to normalize hyperosmolality. In the other patient, however, DDAVP decreased the serum sodium level even with small doses.

L22 ANSWER 21 OF 21 CAPLUS COPYRIGHT 2006 ACS on STN

1966:21723 Document No. 64:21723 Original Reference No. 64:4041h,4042a-c The origin of antidiuretic substances and an explanation of the hypernatraemia in the Schwartz-Bartter syndrome. Lebacqz, E.; Delaere, J. (Univ. Louvain, Belg.). Annales d'Endocrinologie, 26(3), 375-82 (French) 1965. CODEN: ANENAG. ISSN: 0003-4266.

AB Two patients showed the metabolic disturbances of the Schwartz-Bartter syndrome (Schwartz, et al., Am. J. Med. 23, 529-42(1957)): hypernatraemia and urine hyperosmolality, despite hyponatremia and low plasma osmolality. The 1st patient had a duodenal cancer (118 g.), and the natriuria was not modified by the administration of aldosterone (200  $\gamma$ /day for 2 days). The condition was not modified by the oral administration of 100 ml. of 50% EtOH, nor by a perfusion of 120 milliunits of vasopressin over 1 hr. Exts. of urine and of the duodenal tumor were tested for arginine vasopressin. A 24-hr. urine sample contained 1200 milliunits of vasopressin, and the tumor (11 g. of Me<sub>2</sub>CO powder) contained 8 milliunits/mg. of powder. The antidiuretic activity of the exts. disappeared on treatment with thioglycolate. The 2nd patient had a cancer of the bronchus, and whereas restriction of H<sub>2</sub>O intake ameliorated the condition, treatment with cortisone, prednisone, 95% EtOH, diphenylhydantoin, hygroton, or 20% mannitol had no effect. The natriuresis was due mainly to a disturbed proximal reabsorption of Na<sup>+</sup>, and not to an inhibition of aldosterone secretion or an increased glomerular filtration rate in these patients. 34 references.

=> s (ogata e?/au or onuma e?/au or tsunenari t?/au or saito h?/au or azuma y?/au)  
L23 36476 (OGATA E?/AU OR ONUMA E?/AU OR TSUNENARI T?/AU OR SAITO H?/AU  
OR AZUMA Y?/AU)

=> s L23 and PTHrP antibod?  
L24 19 L23 AND PTHRP ANTIBOD?

=> dup remove L24  
PROCESSING COMPLETED FOR L24  
L25 8 DUP REMOVE L24 (11 DUPLICATES REMOVED)

=> d L25 1-8 cbib abs

L25 ANSWER 1 OF 8 MEDLINE on STN

DUPLICATE 1

2005285209. PubMed ID: 15930357. Increased renal calcium reabsorption by parathyroid hormone-related protein is a causative factor in the development of humoral hypercalcemia of malignancy refractory to osteoclastic bone resorption inhibitors. Onuma Etsuro; Azuma Yumiko; Saito Hidemi; Tsunenari Toshiaki; Watanabe Toshihiko; Hirabayashi Manabu; Sato Koh; Yamada-Okabe Hisafumi; Ogata Etsuro. (Pharmaceutical Department IV, Chugai Research Laboratories, Chugai Pharmaceutical, Co., Ltd., Kanagawa, Japan.) Clinical cancer research : an official journal of the American Association for Cancer Research, (2005 Jun 1) Vol. 11, No. 11, pp. 4198-203. Journal code: 9502500. ISSN: 1078-0432. Pub. country: United States. Language: English.

AB PURPOSE: Bisphosphonate and calcitonin lower blood calcium in humoral hypercalcemia of malignancy (HHM) by suppressing osteoclastic bone resorption, but repeated administration of these drugs often leads to relapse. In this study, we examined the roles of parathyroid hormone-related protein (PTHrP) in the development of bisphosphonate- and calcitonin-refractory HHM. EXPERIMENTAL DESIGN: Nude rats bearing the LC-6 JCK tumor xenograft (LC-6 rats) exhibited high bone turnover and HHM. Repeated administration of alendronate induced a sustained suppression of the bone resorption, but it caused only early and transient reduction of the blood calcium levels, leading to unresponsiveness to the drug. Because high blood levels of PTHrP were detected in the LC-6 rats, those that developed alendronate-refractory HHM were treated with an anti-PTHrP antibody. RESULTS: Administration of anti-PTHrP antibody to animals that received repeated administration of alendronate, thereby developing alendronate-refractory HHM, resulted in an increase in fractional excretion of calcium and a marked decrease of blood calcium level. Drug-refractory HHM was also observed in animals that received another osteoclast inhibitor, an eel calcitonin analogue elcatonin. The blood calcium level decreased after the initial administration of elcatonin, but it eventually became elevated during repeated administration. Administration of the anti-PTHrP antibody, but not of alendronate, effectively reduced the blood calcium of the animals that developed elcatonin-refractory HHM. CONCLUSION: High levels of circulating PTHrP and the resulting augmentation of renal calcium reabsorption is one of the major causes of the emergence of osteoclast inhibitor-refractory HHM. Thus, blockage of PTHrP functions by a neutralizing antibody against PTHrP would benefit patients who develop bisphosphonate- or calcitonin-refractory HHM.

L25 ANSWER 2 OF 8 MEDLINE on STN DUPLICATE 2  
2005311286. PubMed ID: 15800941. Parathyroid hormone-related protein (PTHrP) as a causative factor of cancer-associated wasting: possible involvement of PTHrP in the repression of locomotor activity in rats bearing human tumor xenografts. Onuma Etsuro; Tsunenari Toshiaki; Saito Hidemi; Sato Koh; Yamada-Okabe Hisafumi; Ogata Etsuro. (Pharmaceutical Research Department IV, Kamakura Research Laboratories, Chugai Pharmaceutical Co., Kanagawa, Japan.) International journal of cancer. Journal international du cancer, (2005 Sep 1) Vol. 116, No. 3, pp. 471-8. Journal code: 0042124. ISSN: 0020-7136. Pub. country: United States. Language: English.

AB Nude rats bearing the LC-6-JCK human lung cancer xenograft displayed cancer-associated wasting syndrome in addition to humoral hypercalcemia of malignancy. In these rats, not only PTHrP but also several other human proinflammatory cytokines, such as IL-6, leukemia-inducing factor, IL-8, IL-5 and IL-11, were secreted to the bloodstream. Proinflammatory cytokines induce acute-phase reactions, as evidenced by a decrease of serum albumin and an increase in alpha<sub>2</sub>-acid glycoprotein. Tumor resection abolished the production of proinflammatory cytokines and improved acute-phase reactions, whereas anti-PTHrP antibody affected neither proinflammatory cytokine production nor

acute-phase reactions. Nevertheless, tumor resection and administration of anti-PTHrP antibody similarly and markedly attenuated not only hypercalcemia but also loss of fat, muscle and body weight. Body weight gain by anti-PTHrP antibody was associated with increased food consumption; increased body weight from anti-PTHrP antibody was observed when animals were freely fed but not when they were given the same feeding as those that received only vehicle. Furthermore, nude rats bearing LC-6-JCK showed reduced locomotor activity, less eating and drinking and low blood phosphorus; and anti-PTHrP antibody restored them. Although alendronate, a bisphosphonate drug, decreased blood calcium, it affected neither locomotor activity nor serum phosphorus level. These results indicate that PTHrP represses physical activity and energy metabolism independently of hypercalcemia and proinflammatory cytokine actions and that deregulation of such physiologic activities and functions by PTHrP is at least in part involved in PTHrP-induced wasting syndrome.

L25 ANSWER 3 OF 8 MEDLINE on STN DUPLICATE 3  
2003534792. PubMed ID: 14613038. Treatment of malignancy-associated hypercalcemia and cachexia with humanized anti-parathyroid hormone-related protein antibody. Sato Koh; Onuma Etsuro; Yocum Richard C; Ogata Etsuro. (Department of International Coordination, Chugai Pharmaceutical Co, Ltd, Skizuuoka, Japan. ) Seminars in oncology, (2003 Oct) Vol. 30, No. 5 Suppl 16, pp. 167-73. Ref: 11. Journal code: 0420432. ISSN: 0093-7754. Pub. country: United States. Language: English.

AB Parathyroid hormone-related protein (PTHrP) plays a central role in humoral hypercalcemia of malignancy (HHM), which is one of the most frequent paraneoplastic syndromes. PTHrP produced by the tumor acts through a common PTH/PTHrP receptor to promote bone resorption, inhibit calcium excretion from the kidney, and induce hypercalcemia. Patients with HHM often develop cachexia associated with typical symptoms such as anorexia, malaise, nausea, constipation, polyuria, polydipsia, and confusion. The etiology of the cachexia is not fully understood but is thought to be caused by hypercalcemia and various cytokines such as interleukin-6, tumor necrosis factor- $\alpha$ , leukemia inhibitory factor, and others. In this study, we investigated the role of PTHrP in hypercalcemia and cachexia in HHM by using humanized anti-PTHrP antibody. A mouse monoclonal antibody that binds to PTHrP amino acid sequence 1-34 and inhibits PTHrP function has been humanized to create a specific and potent agent for the treatment of patients with HHM. The mouse monoclonal antibody has been shown to have antihypercalcemic activity against nude mice bearing human tumors. Because a mouse antibody is highly immunogenic in human patients, the complementarity-determining regions from the mouse antibody were grafted into a human antibody. The resulting humanized antibody specifically recognizes PTHrP(1-34) and neutralizes PTHrP functions in vitro and in vivo. The humanized anti-PTHrP antibody was administered intravenously to HHM model animals bearing tumors such as LC-6 human lung carcinoma. These animals showed symptoms similar to those of patients with HHM (eg, hypercalcemia and cachexia). The humanized anti-PTHrP antibody-treated animals responded with normalization of blood ionized calcium level through an improvement of bone metabolism and calcium excretion. Moreover, the treated animals also showed an improvement in body weight, ultramotivity, metabolic alkalosis, food consumption, water intake, serum phosphorus, and renal function. Consequently, the humanized antibody-treated animals experienced complete resolution of hypercalcemia and cachexia. These results suggest that the humanized antibody would be an effective and beneficial agent for patients with HHM, and that PTHrP is a major pathogenetic factor of hypercalcemia and cachexia in patients with HHM.

L25 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

2002:888597 Document No. 138:3671 Angiogenesis inhibitors that block binding of PTH-related peptide to its receptor for use as antitumor agents.

Saito, Hidemi; Tsunenari, Toshiaki; Onuma, Etsuro; Kato, Atsuhiko; Suzuki, Masami (Chugai Seiyaku Kabushiki Kaisha, Japan). PCT Int. Appl. WO 2002092133 A1 20021121, 110 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (Japanese). CODEN: PIXXD2.

AB APPLICATION: WO 2002-JP4586 20020510. PRIORITY: JP 2001-140659 20010510. It is found out that angiogenesis can be inhibited by a substance which inhibits the binding of a parathyroid hormone-associated peptide (e.g. PTHrP) to its receptor. The angiogenesis inhibitors can be anti-PTHrP antibodies, antibody fragments, humanized or chimeric antibodies, PTH receptor antagonists, or antisense oligonucleotides specific to PTHrP. These modified anti-PTHrP antibodies and PTH receptor antagonists are useful as antitumor agents and bone metastasis inhibitors.

L25 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

2001:31355 Document No. 134:99582 Remedies for drug-resistant hypercalcemia.

Saito, Hidemi; Tsunenari, Toshiaki; Onuma, Etsuro (Chugai Seiyaku Kabushiki Kaisha, Japan). PCT Int. Appl. WO 2001002012 A1 20010111, 118 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (Japanese). CODEN: PIXXD2. APPLICATION: WO 2000-JP4523 20000706. PRIORITY: JP 1999-192270 19990706.

AB Remedies for drug-resistant hypercalcemia which contain as the active ingredient a substance inhibiting the binding of a parathyroid hormone-related peptide to its receptor. Therapeutics for drug-resistant hypercalcemia include bone resorption inhibitor (e.g. bisphosphates and/or calcitonin), calcium excretion promoter, intestinal calcium absorption inhibitor, or loop diuretic. The PTHrP and receptor-binding inhibitors are PTHrP receptor antagonist such as anti-PTHrP antibodies or fragments or chimeric antibodies.

L25 ANSWER 6 OF 8 MEDLINE on STN

DUPLICATE 4

2000354698. PubMed ID: 10898333. Parathyroid hormone-related protein as a potential target of therapy for cancer-associated morbidity. Ogata E. (Japanese Foundation for Cancer Research, Tokyo. ) Cancer, (2000 Jun 15) Vol. 88, No. 12 Suppl, pp. 2909-11. Journal code: 0374236. ISSN: 0008-543X. Pub. country: United States. Language: English.

AB BACKGROUND: Proinflammatory cytokines are involved in the genesis of cancer-associated cachexia. Parathyroid hormone-related protein (PTHrP) is the causative agent in humoral hypercalcemia of malignancy (HHM) and is frequently secreted from various kinds of solid tumors as well as from adult T-cell leukemia/lymphoma. PTHrP, like PTH, acts on PTH receptor type 1 (PTH1R). Activation of PTH1R may lead to stimulation of secretion of proinflammatory cytokines. It is expected, therefore, that PTHrP constitutes a key factor in the activation of the proinflammatory and cachectogenic cytokine network and consequently in the development of cachexia in patients with cancer. METHODS: Two groups of cancer-bearing patients of similar clinical backgrounds were enrolled. Plasma

concentrations of PTHrP and cytokines were measured with immunoradiometric assay and radioimmunoassay, respectively. Cancer tissues from patients with HHM were transplanted into nude mice or nude rats. The effects of humanized antihuman PTHrP antibody were examined.

RESULTS: In clinical studies, Group B patients (with elevated plasma PTHrP), compared with Group A patients (with normal plasma PTHrP), tended to exhibit higher plasma levels of tumor necrosis factor (TNF)-alpha (P = 0.13), interleukin (IL)-5 (P = 0.08), and IL-8 (P = 0.08), and had significantly higher levels of IL-6 (P = < or = 0.01). The levels of TNF-alpha and IL-6 correlated with those of PTHrP. In animal studies, the antibody caused a prompt and sustained decline in serum calcium. This response was accompanied by improvements in food intake, drinking, body weight gain, and general behavior. It also ameliorated the suppression of serum ADH. When those effects were compared with those induced either by bisphosphonate or calcitonin, it turned out that not all of the beneficial effects of the antibody were directly correlated with the depression of blood calcium. CONCLUSIONS: PTHrP is a promising molecular target for the development of a novel mode of treatment for patients with cancer-associated morbidity.

L25 ANSWER 7 OF 8 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN 2000:244979 Document No.: PREV200000244979. The possibility of utilizing humanized anti-PTHrP antibody as an anti-HHM/cachexia agent. Onuma, Etsuro [Reprint author]; Saito, H.; Azuma, Y.; Shimizu, N.; Tsunenari, T.; Sato, K.; Ogata, E.. Chugai Pharmaceutical, Shizuoka, Japan. Proceedings of the American Association for Cancer Research Annual Meeting, (March, 2000) No. 41, pp. 287. print. Meeting Info.: 91st Annual Meeting of the American Association for Cancer Research. San Francisco, California, USA. April 01-05, 2000. ISSN: 0197-016X. Language: English.

L25 ANSWER 8 OF 8 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN 2000:534320 Document No.: PREV200000534320. PTHrP: The most potent cachectogenic factor shown in an animal model. Ogata, E. [Reprint author]; Takahashi, S. [Reprint author]; Onuma, E.; Sato, K.. Cancer Institute Hospital, Tokyo, Japan. Bone (New York), (October, 2000) Vol. 27, No. 4 Supplement, pp. 21S. print. Meeting Info.: 2000 International Bone and Hormone Meeting. Hamilton Island, Great Barrier Reef, Queensland, Australia. November 04-07, 2000. International Bone and Mineral Society. CODEN: BONEDL. ISSN: 8756-3282. Language: English.

=> s 123 and treatment  
L26 4243 L23 AND TREATMENT  
  
=> s 126 and vasopressin level  
L27 0 L26 AND VASOPRESSIN LEVEL  
  
=> s 126 and vesopressin  
L28 0 L26 AND VESOPRESSIN  
  
=> s 126 and hyperosmolarity  
L29 0 L26 AND HYPEROSMOLARITY  
  
=> s 126 and anti-PTHrP  
L30 20 L26 AND ANTI-PTHRP

=> dup remove l30  
PROCESSING COMPLETED FOR L30  
L31 6 DUP REMOVE L30 (14 DUPLICATES REMOVED)



=> d 131 1-6 cbib abs

L31 ANSWER 1 OF 6 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

2005255152 EMBASE Increased renal calcium reabsorption by parathyroid hormone-related protein is a causative factor in the development of humoral hypercalcemia of malignancy refractory to osteoclastic bone resorption inhibitors. Onuma E.; Azuma Y.; Saito H.; Tsunenari T.; Watanabe T.; Hirabayashi M.; Sato K.; Yamada-Okabe H.; Ogata E.. H. Yamada-Okabe, Pharmaceutical Research Department IV, Kamakura Research Laboratories, Chugai Pharmaceutical Co., Ltd., 200 Kajiwarra, Kamakura, 247-8530 Kanagawa, Japan. okabehsf@chugai-pharm.co.jp. Clinical Cancer Research Vol. 11, No. 11, pp. 4198-4203 1 Jun 2005.

Refs: 18.

ISSN: 1078-0432. CODEN: CCREF4

Pub. Country: United States. Language: English. Summary Language: English. Entered STN: 20050707. Last Updated on STN: 20050707

AB Purpose: Bisphosphonate and calcitonin lower blood calcium in humoral hypercalcemia of malignancy (HHM) by suppressing osteoclastic bone resorption, but repeated administration of these drugs often leads to relapse. In this study, we examined the roles of parathyroid hormone-related protein (PTHrP) in the development of bisphosphonate- and calcitonin-refractory HHM. Experimental Design: Nude rats bearing the LC-6 JCK tumor xenograft (LC-6 rats) exhibited high bone turnover and HHM. Repeated administration of alendronate induced a sustained suppression of the bone resorption, but it caused only early and transient reduction of the blood calcium levels, leading to unresponsiveness to the drug. Because high blood levels of PTHrP were detected in the LC-6 rats, those that developed alendronate-refractory HHM were treated with an anti-PTHrP antibody. Results: Administration of anti-PTHrP antibody to animals that received repeated administration of alendronate, thereby developing alendronate-refractory HHM, resulted in an increase in fractional excretion of calcium and a marked decrease of blood calcium level. Drug-refractory HHM was also observed in animals that received another osteoclast inhibitor, an eel calcitonin analogue elcatonin. The blood calcium level decreased after the initial administration of elcatonin, but it eventually became elevated during repeated administration. Administration of the anti-PTHrP antibody, but not of alendronate, effectively reduced the blood calcium of the animals that developed elcatonin-refractory HHM. Conclusion: High levels of circulating PTHrP and the resulting augmentation of renal calcium reabsorption is one of the major causes of the emergence of osteoclast inhibitor-refractory HHM. Thus, blockage of PTHrP functions by a neutralizing antibody against PTHrP would benefit patients who develop bisphosphonate- or calcitonin-refractory HHM. .COPYRG. 2005 American Association for Cancer Research.

L31 ANSWER 2 OF 6 MEDLINE on STN DUPLICATE 1

2005628438. PubMed ID: 16309168. Humanized monoclonal antibody against parathyroid hormone-related protein suppresses osteolytic bone metastasis of human breast cancer cells derived from MDA-MB-231. Saito Hidemi; Tsunenari Toshiaki; Onuma Etsuro; Sato Koh; Ogata Etsuro; Yamada-Okabe Hisafumi. (Pharmaceutical Research Department III, Kamakura Research Laboratories, Chugai Pharmaceutical Co., Ltd., Kamakura, Kanagawa 247-8530, Japan. ) Anticancer research, (2005 Nov-Dec) Vol. 25, No. 6B, pp. 3817-23. Journal code: 8102988. ISSN: 0250-7005. Pub. country: Greece. Language: English.

AB BACKGROUND: Parathyroid hormone-related protein (PTHrP) has been implicated in bone metastasis. However, the effects on bone metastasis of blocking the PTHrP function have not been tested in the clinic. Here, the

effects of a humanized anti-PTHrP monoclonal antibody (mAb) on bone metastasis in a human xenograft model are shown. MATERIALS AND METHODS: Subline MDA-5a, with high bone metastatic activity, was established from the human breast cancer cell line MDA-MB-231. Mice were injected with MDA-5a and an anti-PTHrP monoclonal antibody (mAb) raised against human PTHrP (1-34); bone metastasis was evaluated by X-ray photography. RESULTS: MDA-5a produced elevated levels of PTHrP, Interleukin 8 (IL-8), IL-6 and matrix metalloproteinase 1 (MMP-1) and frequently metastasized to the bone. Administration of the humanized anti-PTHrP mAb significantly suppressed osteolytic bone metastasis of MDA-5a and caused osteogenesis at the sites of metastasis. CONCLUSION: The humanized anti-PTHrP mAb was effective against bone metastasis by inducing osteogenesis and, therefore, will provide a new treatment option for bone metastasis in breast cancer.

L31 ANSWER 3 OF 6 MEDLINE on STN DUPLICATE 2  
2004546213. PubMed ID: 15517871. Generation of a humanized monoclonal antibody against human parathyroid hormone-related protein and its efficacy against humoral hypercalcemia of malignancy. Onuma Etsuro ; Sato Koh; Saito Hidemi; Tsunenari Toshiaki; Ishii Kimie; Esaki Keiko; Yabuta Naohiro; Wakahara Yuji; Yamada-Okabe Hisafumi; Ogata Etsuro. (Chugai Research Laboratories, Chugai Pharmaceutical Co. Ltd., 200 Kajiwara, Kamakura, Kanagawa, Japan. ) Anticancer research, (2004 Sep-Oct) Vol. 24, No. 5A, pp. 2665-73. Journal code: 8102988. ISSN: 0250-7005. Pub. country: Greece. Language: English.

AB A humanized monoclonal antibody against parathyroid hormone-related protein (PTHrP) was generated from the mouse monoclonal antibody raised against the peptide corresponding to the N-terminal 34 amino acids of the human PTHrP [(PTHrP(1-34))]. The humanized antibody interacted with the PTHrP(1-34) with a K<sub>d</sub> value of 1.90 x 10<sup>-10</sup> M, and the epitope resides between the amino acids 20 and 30 of the PTHrP. PTHrP(1-34) significantly increased the intracellular cAMP levels in the rat osteosarcoma cells that expressed PTHrP, and the 5 microg/mL or higher concentrations of the humanized antibody almost completely blocked the PTHrP-induced cAMP production even in the presence of 2 microg/mL PTHrP(1-34), demonstrating its ability to fully neutralize PTHrP function. There was no significant difference in the potency of the mouse, chimera, or the humanized antibodies to suppress the PTHrP-induced increase in the intracellular cAMP in ROS cells. Furthermore, at the same doses, the administration of the chimera or the humanized antibody was equally effective in reducing the blood ionized calcium levels of hypercalcemic mice bearing the PAN-7-JCK human pancreatic cancer xenograft or the LC-6-JCK human lung cancer xenograft that secreted PTHrP. Thus, humanized anti-PTHrP may be useful for the treatment of the humoral hypercalcemia of malignancy in humans.

L31 ANSWER 4 OF 6 MEDLINE on STN DUPLICATE 3  
2003534792. PubMed ID: 14613038. Treatment of malignancy-associated hypercalcemia and cachexia with humanized anti-parathyroid hormone-related protein antibody. Sato Koh; Onuma Etsuro; Yocum Richard C; Ogata Etsuro. (Department of International Coordination, Chugai Pharmaceutical Co, Ltd, Skizuuoka, Japan. ) Seminars in oncology, (2003 Oct) Vol. 30, No. 5 Suppl 16, pp. 167-73. Ref: 11. Journal code: 0420432. ISSN: 0093-7754. Pub. country: United States. Language: English.

AB Parathyroid hormone-related protein (PTHrP) plays a central role in humoral hypercalcemia of malignancy (HHM), which is one of the most frequent paraneoplastic syndromes. PTHrP produced by the tumor acts through a common PTH/PTHrP receptor to promote bone resorption, inhibit calcium excretion from the kidney, and induce hypercalcemia. Patients with HHM often develop cachexia associated with typical symptoms such as

anorexia, malaise, nausea, constipation, polyuria, polydipsia, and confusion. The etiology of the cachexia is not fully understood but is thought to be caused by hypercalcemia and various cytokines such as interleukin-6, tumor necrosis factor- $\alpha$ , leukemia inhibitory factor, and others. In this study, we investigated the role of PTHrP in hypercalcemia and cachexia in HHM by using humanized anti-PTHrP antibody. A mouse monoclonal antibody that binds to PTHrP amino acid sequence 1-34 and inhibits PTHrP function has been humanized to create a specific and potent agent for the treatment of patients with HHM. The mouse monoclonal antibody has been shown to have antihypercalcemic activity against nude mice bearing human tumors. Because a mouse antibody is highly immunogenic in human patients, the complementarity-determining regions from the mouse antibody were grafted into a human antibody. The resulting humanized antibody specifically recognizes PTHrP(1-34) and neutralizes PTHrP functions in vitro and in vivo. The humanized anti-PTHrP antibody was administered intravenously to HHM model animals bearing tumors such as LC-6 human lung carcinoma. These animals showed symptoms similar to those of patients with HHM (eg, hypercalcemia and cachexia). The humanized anti-PTHrP antibody-treated animals responded with normalization of blood ionized calcium level through an improvement of bone metabolism and calcium excretion. Moreover, the treated animals also showed an improvement in body weight, ulromotivity, metabolic alkalosis, food consumption, water intake, serum phosphorus, and renal function. Consequently, the humanized antibody-treated animals experienced complete resolution of hypercalcemia and cachexia. These results suggest that the humanized antibody would be an effective and beneficial agent for patients with HHM, and that PTHrP is a major pathogenetic factor of hypercalcemia and cachexia in patients with HHM.

- L31 ANSWER 5 OF 6 MEDLINE on STN DUPLICATE 4  
 2003428817. PubMed ID: 12969787. Monoclonal antibody to parathyroid hormone-related protein induces differentiation and apoptosis of chondrosarcoma cells. Miyaji Takahiro; Nakase Takano; Onuma Eturo; Sato Koh; Myoui Akira; Tomita Tetsuya; Joyama Susumu; Ariga Kenta; Hashimoto Jun; Ueda Takafumi; Yoshikawa Hideki. (Department of Orthopaedic Surgery, Osaka University Medical School, 2-2 Yamadaoka, Suita 565-0871, Japan.. miyaji@ort.med.osaka-u.ac.jp) . Cancer letters, (2003 Sep 25) Vol. 199, No. 2, pp. 147-55. Journal code: 7600053. ISSN: 0304-3835. Pub. country: Ireland. Language: English.
- AB We investigated the effects of treatment with anti-parathyroid hormone-related protein (1-34) monoclonal murine antibody (anti-PTHrP MoAb) on apoptosis and the differentiation of chondrosarcoma HTB-94 cells. Treatment with anti-PTHrP MoAb accelerated apoptosis of HTB-94 cells in a dose-dependent manner, and anti-PTHrP MoAb also promoted the chondrogenic differentiation of HTB-94 cells. The induction of apoptosis by anti-PTHrP MoAb via imbalance of Bcl-2/Bax ratio and activation of caspase-3 may provide a mechanistic explanation for its potential antitumor effects. Our results suggest the possibility that anti-PTHrP MoAb may be beneficial as a new treatment for chondrosarcoma.

- L31 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2006 ACS on STN  
 2001:31355 Document No. 134:99582 Remedies for drug-resistant hypercalcemia. Saito, Hidemi; Tsunenari, Toshiaki; Onuma, Etsuro (Chugai Seiyaku Kabushiki Kaisha, Japan). PCT Int. Appl. WO 200102012 A1 20010111, 118 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MG, MK, MN, MW, MX, MY, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,

UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (Japanese). CODEN: PIXXD2. APPLICATION: WO 2000-JP4523 20000706. PRIORITY: JP 1999-192270 19990706.

AB Remedies for drug-resistant hypercalcemia which contain as the active ingredient a substance inhibiting the binding of a parathyroid hormone-related peptide to its receptor. Therapeutics for drug-resistant hypercalcemia include bone resorption inhibitor (e.g. bisphosphates and/or calcitonin), calcium excretion promoter, intestinal calcium absorption inhibitor, or loop diuretic. The PTHrP and receptor-binding inhibitors are PTHrP receptor antagonist such as anti-PTHrP antibodies or fragments or chimeric antibodies.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	201.88	202.09
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-5.25	-5.25

STN INTERNATIONAL LOGOFF AT 10:21:23 ON 27 DEC 2006

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSSPTA1644PNH

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS	3	AUG 06	FSTA enhanced with new thesaurus edition
NEWS	4	AUG 13	CA/Caplus enhanced with additional kind codes for granted patents
NEWS	5	AUG 20	CA/Caplus enhanced with CAS indexing in pre-1907 records
NEWS	6	AUG 27	Full-text patent databases enhanced with predefined patent family display formats from INPADOCDB

NEWS 7 AUG 27 USPATOLD now available on STN  
 NEWS 8 AUG 28 CAS REGISTRY enhanced with additional experimental  
 spectral property data  
 NEWS 9 SEP 07 STN AnaVist, Version 2.0, now available with Derwent  
 World Patents Index  
 NEWS 10 SEP 13 FORIS renamed to SOPIS  
 NEWS 11 SEP 13 INPADOCDB enhanced with monthly SDI frequency  
 NEWS 12 SEP 17 CA/CAPLUS enhanced with printed CA page images from  
 1967-1998  
 NEWS 13 SEP 17 CAPLUS coverage extended to include traditional medicine  
 patents  
 NEWS 14 SEP 24 EMBASE, EMBAL, and LEMBASE reloaded with enhancements  
 NEWS 15 OCT 02 CA/CAPLUS enhanced with pre-1907 records from Chemisches  
 Zentralblatt  
 NEWS 16 OCT 19 BEILSTEIN updated with new compounds  
 NEWS 17 NOV 15 Derwent Indian patent publication number format enhanced  
 NEWS 18 NOV 19 WPIX enhanced with XML display format  
 NEWS 19 NOV 30 ICSD reloaded with enhancements  
 NEWS 20 DEC 04 LINPADOCDB now available on STN  
 NEWS 21 DEC 14 BEILSTEIN pricing structure to change  
 NEWS 22 DEC 17 USPATOLD added to additional database clusters  
 NEWS 23 DEC 17 IMSDRUGCONF removed from database clusters and STN  
 NEWS 24 DEC 17 DGENE now includes more than 10 million sequences  
 NEWS 25 DEC 17 TOXCENTER enhanced with 2008 MeSH vocabulary in  
 MEDLINE segment  
 NEWS 26 DEC 17 MEDLINE and LEMBASE updated with 2008 MeSH vocabulary  
 NEWS 27 DEC 17 CA/CAPLUS enhanced with new custom IPC display formats  
 NEWS 28 DEC 17 STN Viewer enhanced with full-text patent content  
 from USPATOLD  
 NEWS 29 JAN 02 STN pricing information for 2008 now available  
 NEWS 30 JAN 16 CAS patent coverage enhanced to include exemplified  
 prophetic substances  
  
 NEWS EXPRESS 19 SEPTEMBER 2007: CURRENT WINDOWS VERSION IS V8.2,  
 CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
 AND CURRENT DISCOVER FILE IS DATED 19 SEPTEMBER 2007.  
  
 NEWS HOURS STN Operating Hours Plus Help Desk Availability  
 NEWS LOGIN Welcome Banner and News Items  
 NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that  
 specific topic.

All use of STN is subject to the provisions of the STN Customer  
 agreement. Please note that this agreement limits use to scientific  
 research. Use for software development or design or implementation  
 of commercial gateways or other similar uses is prohibited and may  
 result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 09:22:42 ON 24 JAN 2008

=> file medline embase biosis scisearch caplus  
 COST IN U.S. DOLLARS

	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 09:23:05 ON 24 JAN 2008

FILE 'EMBASE' ENTERED AT 09:23:05 ON 24 JAN 2008

Copyright (c) 2008 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 09:23:05 ON 24 JAN 2008  
Copyright (c) 2008 The Thomson Corporation

FILE 'SCISEARCH' ENTERED AT 09:23:05 ON 24 JAN 2008  
Copyright (c) 2008 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 09:23:05 ON 24 JAN 2008  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

=> s humanized antibod?  
L1 3157 HUMANIZED ANTIBOD?

=> s l1 and PTHrP  
L2 16 L1 AND PTHRP

=> dup remove l2  
PROCESSING COMPLETED FOR L2  
L3 10 DUP REMOVE L2 (6 DUPLICATES REMOVED)

=> s l3 and pd<20000720  
L4 2 FILES SEARCHED...  
L4 2 L3 AND PD<20000720

=> d l4 cbib abs

L4 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2008 ACS on STN  
2001:30858 Document No. 135:59794 The application of BIA technology in the optimization of humanized mouse anti-parathyroid hormone-related protein ( PTHrP) antibodies. Esaki, Keiko (Pharmaceutical Technology Lab., Chugai Pharmaceutical Co., Ltd., Gotemba, 412-8513, Japan). Real-Time Analysis of Biomolecular Interactions, 195-200. Editor(s): Nagata, Kazuhiro; Handa, Hiroshi. Springer-Verlag Tokyo: Tokyo, Japan. (English) 2000. CODEN: 69AUQB.

AB A review with 5 refs. Topics discussed include the preparation of the sensor chip surface; the interaction between mouse monoclonal antibodies and the immobilized [Cys35]-PTHrP(1-35); optimization of humanized antibodies; and the kinetic anal. of humanized antibodies and the immobilized [Cys35]-PTHrP(1-35).

=> d l4 1-2 cbib abs

L4 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2008 ACS on STN  
2001:30858 Document No. 135:59794 The application of BIA technology in the optimization of humanized mouse anti-parathyroid hormone-related protein ( PTHrP) antibodies. Esaki, Keiko (Pharmaceutical Technology Lab., Chugai Pharmaceutical Co., Ltd., Gotemba, 412-8513, Japan). Real-Time Analysis of Biomolecular Interactions, 195-200. Editor(s): Nagata, Kazuhiro; Handa, Hiroshi. Springer-Verlag Tokyo: Tokyo, Japan. (English) 2000. CODEN: 69AUQB.

AB A review with 5 refs. Topics discussed include the preparation of the sensor chip surface; the interaction between mouse monoclonal antibodies and the immobilized [Cys35]-PTHrP(1-35); optimization of humanized antibodies; and the kinetic anal. of humanized antibodies and the immobilized [Cys35]-PTHrP(1-35).

L4 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2008 ACS on STN

1998:210769 Document No. 128:293971 Antibody against human parathormone related peptides. Sato, Kou; Wakahara, Yuji; Yabuta, Naohiro (Chugai Seiyaku K. K., Japan; Sato, Kou; Wakahara, Yuji; Yabuta, Naohiro). PCT Int. Appl. WO 9813388 A1 19980402, 184 pp. DESIGNATED STATES: W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (Japanese). CODEN: PIXXD2. APPLICATION: WO 1997-JP3382 19970924. PRIORITY: JP 1996-255196 19960926; JP 1997-214168 19970724.

AB An antibody against human parathormone related peptides (hPTHrP), a DNA encoding the antibody, a recombinant vector containing the DNA, a transformant prepared with the vector, a process for producing the antibody, and the application of the antibody. These humanized anti-hPTHrP antibody fragments are useful for treating hypercalcemia accompanying malignancy or hypophosphatemia.

=> s anti-hPTHrP

L5 16 ANTI-HPTHR

=> s l5 and treat?

L6 1 L5 AND TREAT?

=> d l6 cbib abs

L6 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN

1998:210769 Document No. 128:293971 Antibody against human parathormone related peptides. Sato, Kou; Wakahara, Yuji; Yabuta, Naohiro (Chugai Seiyaku K. K., Japan; Sato, Kou; Wakahara, Yuji; Yabuta, Naohiro). PCT Int. Appl. WO 9813388 A1 19980402, 184 pp. DESIGNATED STATES: W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (Japanese). CODEN: PIXXD2. APPLICATION: WO 1997-JP3382 19970924. PRIORITY: JP 1996-255196 19960926; JP 1997-214168 19970724.

AB An antibody against human parathormone related peptides (hPTHrP), a DNA encoding the antibody, a recombinant vector containing the DNA, a transformant prepared with the vector, a process for producing the antibody, and the application of the antibody. These humanized anti-hPTHrP antibody fragments are useful for treating hypercalcemia accompanying malignancy or hypophosphatemia.

=> s treat?

L7 13159831 TREAT?

=> s l7 and hypercalcemia

L8 16983 L7 AND HYPERCALCEMIA

=> s l8 and polyuria

L9 212 L8 AND POLYURIA

=> s l9 and antibod?

L10 10 L9 AND ANTIBOD?

=> s l10 and PTHrP1-34  
L11 0 L10 AND PTHRP1-34

=> s l10 and PTHrP  
L12 4 L10 AND PTHRP

=> dup remove l12  
PROCESSING COMPLETED FOR L12  
L13 2 DUP REMOVE L12 (2 DUPLICATES REMOVED)

=> d l13 1-2 cbib abs

L13 ANSWER 1 OF 2 MEDLINE on STN  
2006057279. PubMed ID: 16444366. [Hypercalcemia of malignancy: clinical features, diagnosis and treatment]. A hipercalcemia nas malignidades: aspectos clinicos, diagnosticos e terapeuticos. Farias Maria Lucia F de. (Faculdade de Medicina, Universidade Federal do Rio de Janeiro, RJ.. fleiuss@hucff.ufrj.br) . Arquivos brasileiros de endocrinologia e metabologia, (2005 Oct) Vol. 49, No. 5, pp. 816-24. Electronic Publication: 2006-01-23. Ref: 64. Journal code: 0403437. ISSN: 0004-2730. Pub. country: Brazil. Language: Portuguese.

AB Hypercalcemia associated with malignancies is reported in up to 20 to 30% of patients with cancer during the course of the disease, and points to a poor prognosis. Symptoms related to the central nervous system, as progressive mental impairment, stupor and coma, predominate. Alterations in kidney function (water-concentrating defect leading to polyuria) and gastrointestinal tract (anorexia, nausea, vomiting) corroborate to dehydration and a further increase in serum calcium. Cancer-induced hypercalcemia may be classified as: 1) local osteolytic hypercalcemia (LOH), due to marked increase in osteoclastic bone resorption in areas surrounding the malignant cells within the marrow space; 2) humoral hypercalcemia of malignancy, caused by the secretion of parathyroid hormone-related protein (PTHrP) by the malignant tumor; 3) ectopic hyperparathyroidism; 4) 1,25(OH)2 D-secreting tumors. Adequate control of hypercalcemia is necessary to give the patient time to respond to anti-cancer therapy. Volume expansion with saline will correct dehydration, improve glomerular filtration and increase urinary calcium excretion, which may be further stimulated by loop diuretics. Intravenous bisphosphonates are the most effective agents to control hypercalcemia, as they block osteoclastic osteolysis and also have antitumoral effects, decreasing bone metastases. New approaches to control the skeletal manifestations of malignancies are anti-PTHrP and anti-RANKL antibodies, osteoprotegerin, and also proteasome inhibitors in the case of multiple myeloma.

L13 ANSWER 2 OF 2 MEDLINE on STN DUPLICATE 1  
2003534792. PubMed ID: 14613038. Treatment of malignancy-associated hypercalcemia and cachexia with humanized anti-parathyroid hormone-related protein antibody. Sato Koh; Onuma Etsuro; Yocum Richard C; Ogata Etsuro. (Department of International Coordination, Chugai Pharmaceutical Co, Ltd, Skizuuoka, Japan. ) Seminars in oncology, (2003 Oct) Vol. 30, No. 5 Suppl 16, pp. 167-73. Ref: 11. Journal code: 0420432. ISSN: 0093-7754. Pub. country: United States. Language: English.

AB Parathyroid hormone-related protein (PTHrP) plays a central role in humoral hypercalcemia of malignancy (HHM), which is one of the most frequent paraneoplastic syndromes. PTHrP produced by the tumor acts through a common PTH/PTHrP receptor to promote bone resorption, inhibit calcium excretion from the kidney, and induce hypercalcemia. Patients with HHM often develop cachexia associated with typical symptoms such as anorexia, malaise, nausea,



constipation, polyuria, polydipsia, and confusion. The etiology of the cachexia is not fully understood but is thought to be caused by hypercalcemia and various cytokines such as interleukin-6, tumor necrosis factor- $\alpha$ , leukemia inhibitory factor, and others. In this study, we investigated the role of PTHrP in hypercalcemia and cachexia in HHM by using humanized anti-PTHrP antibody. A mouse monoclonal antibody that binds to PTHrP amino acid sequence 1-34 and inhibits PTHrP function has been humanized to create a specific and potent agent for the treatment of patients with HHM. The mouse monoclonal antibody has been shown to have antihypercalcemic activity against nude mice bearing human tumors. Because a mouse antibody is highly immunogenic in human patients, the complementarity-determining regions from the mouse antibody were grafted into a human antibody. The resulting humanized antibody specifically recognizes PTHrP(1-34) and neutralizes PTHrP functions in vitro and in vivo. The humanized anti-PTHrP antibody was administered intravenously to HHM model animals bearing tumors such as LC-6 human lung carcinoma. These animals showed symptoms similar to those of patients with HHM (eg, hypercalcemia and cachexia). The humanized anti-PTHrP antibody-treated animals responded with normalization of blood ionized calcium level through an improvement of bone metabolism and calcium excretion. Moreover, the treated animals also showed an improvement in body weight, ultramotivity, metabolic alkalosis, food consumption, water intake, serum phosphorus, and renal function. Consequently, the humanized antibody-treated animals experienced complete resolution of hypercalcemia and cachexia. These results suggest that the humanized antibody would be an effective and beneficial agent for patients with HHM, and that PTHrP is a major pathogenetic factor of hypercalcemia and cachexia in patients with HHM.

```
=> s vasopressin level
L14      3234 VASOPRESSIN LEVEL
```

```
=> s l14 and brain
L15      518 L14 AND BRAIN
```

```
=> s l15 and maintain
L16      13 L15 AND MAINTAIN
```

```
=> dup remove l16
PROCESSING COMPLETED FOR L16
L17      8 DUP REMOVE L16 (5 DUPLICATES REMOVED)
```

```
=> d l17 1-8 cbib abs
```

```
L17  ANSWER 1 OF 8 EMBASE  COPYRIGHT (c) 2008 Elsevier B.V. All rights
      reserved on STN
2004138192 EMBASE  New additions to the intensive care armamentarium. Rice
      T.W.; Bernard G.R.. Dr. T.W. Rice, Div. Allergy, Pulmon./Crit. Care M.,
      Vanderbilt Univ. School of Medicine, Center for Lung Research, Nashville,
      TN 37232-2650, United States. todd.rice@vanderbilt.edu. Drugs of Today
      Vol. 40, No. 2, pp. 157-170 Feb 2004.
      Refs: 107.
      ISSN: 0025-7656. CODEN: MDACAP
      Pub. Country: Spain. Language: English. Summary Language: English.
      Entered STN: 20040415. Last Updated on STN: 20040415
AB   Many advances have improved the care of critically ill patients, but only
      a few have been through the use of pharmaceutical agents. Recently, the
```

US Food and Drug Administration (FDA) approved drotrecogin alfa (activated), or recombinant human activated protein C, for the treatment of patients with a high risk of death from severe sepsis. Drotrecogin alfa (activated) has antiinflammatory, antithrombotic and fibrinolytic properties. When given as a continuous intravenous infusion, recombinant human activated protein C decreases absolute mortality of severely septic patients by 6.1%, resulting in a 19.4% relative reduction in mortality. The absolute reduction in mortality increases to 13% if the population treated is restricted to patients with an APACHE II score greater than 24, as suggested by the FDA. The most frequent and serious side effect is bleeding. Severe bleeds increased from 2% in patients given placebo to 3.5% in patients receiving drotrecogin alfa (activated). The risk of bleeding was only increased during the actual infusion time of the drug, and the bleeding risk returned to placebo levels 24 hours after the infusion was discontinued. Patients treated in the intensive care unit frequently develop anemia, usually severe enough to require at least one transfusion of red blood cells. With the recent discovery of the harmful effects of allogeneic red blood cell transfusions and the increasing shortage of available red blood cell products, emphasis has been placed on minimizing transfusions. Patients who receive exogenous recombinant human erythropoietin maintain higher hemoglobin levels, in spite of requiring fewer transfusions during their stay in the intensive care unit. Recombinant human erythropoietin appears to be effective whether it is given as 300 units/kg of body weight subcutaneously every other day or as 40,000 units subcutaneously every week. Differences in hemoglobin values were not apparent until at least one week of therapy, but they continued to diverge after that initial week. Furthermore, the incidence of adverse events was similar to that of patients receiving placebo and there was no difference in mortality, suggesting that avoidance of blood transfusions did not translate into increased survival. Thus, recombinant human erythropoietin appears to be both safe and effective in treating the anemia found in critically ill patients, but it is less clear that such treatment is cost effective, especially in the higher dose regimens. Hypotension in patients with septic shock is often difficult to correct. Despite enormous dosages of catecholamines, many of these patients continue to have inadequate blood pressures. Inadequate levels of vasopressin have been identified in patients with septic shock, as well as in other patients with hypotension secondary to refractory vasodilatation. Vasopressin is a peptide hormone secreted from the posterior pituitary in response to hyperosmolality, hypovolemia or hypotension. Levels of vasopressin initially rise in patients with septic shock, but as hypotension persists, vasopressin levels fall below normal. Administration of exogenous vasopressin in physiologic dosages significantly increases blood pressure in patients with shock associated with sepsis and other vasodilatory states. This rise in blood pressure is often significant enough that endogenous catecholamines can be decreased and frequently discontinued entirely. Early withdrawal of the vasopressin replacement infusion results in recurrent hypotension. Unfortunately, randomized, blinded, placebo-controlled trials showing improvement in long-term outcomes such as mortality and length of stay are still lacking. .COPYRGHT. 2004 Prous Science. All rights reserved.

- L17 ANSWER 2 OF 8 MEDLINE on STN DUPLICATE 1  
 1998278559. PubMed ID: 9617997. Nitric oxide control of drinking, vasopressin and oxytocin release and blood pressure in dehydrated rats. Liu H; Terrell M L; Bui V; Summy-Long J Y; Kadekaro M. (Division of Neurosurgery, the University of Texas Medical Branch at Galveston, 77555-0517, USA. ) Physiology & behavior, (1998 Mar) Vol. 63, No. 5, pp. 763-9. Journal code: 0151504. ISSN: 0031-9384. Pub. country: United States. Language: English.
- AB Intracerebroventricular (i.c.v.) injection of the inhibitor of NO synthase (NOS), N(G)-nitro-L-arginine methyl ester (L-NAME) (250 microg/5 microl)

attenuated the drinking response in rats deprived of water for 24 h. Moreover, oxytocin (OT) levels in plasma increased after 2 min, whereas both oxytocin and vasopressin levels were elevated at 120 min after intracerebroventricular injection. The delayed effect of L-NAME on both hormones was not observed in dehydrated animals allowed to drink water. Blood pressure remained stable after injection of artificial cerebrospinal fluid (aCSF) in dehydrated rats not allowed to drink. In rats having access to water, however, there was an immediate but transient pressor response (0-5 min) with a delayed hypotension from 45 to 120 min. L-NAME consistently increased blood pressure in a biphasic mode, whether the animals drank or not, with an early peak at 5 min that decayed after 15-30 min and a second pressor response beginning at 30-45 min and remaining elevated at 120 min when the experiment ended. These pressor responses were independent of the adrenal glands. Thus, centrally produced nitric oxide facilitates drinking, inhibits release of vasopressin and oxytocin from the magnocellular system, and maintains resting arterial blood pressure in normally hydrated and dehydrated rats.

L17 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN

1994:103885 Document No. 120:103885 Effect of hypoxemia on the cardiovascular response to intracranial hypertension in postnatal lambs. Kearney, Marguerite L.; Backofen, Joanne E.; Koehler, Raymond C.; Jones, M. Douglas, Jr.; Traystman, Richard J. (Dep. Anesthesiol., Johns Hopkins Med. Inst., Baltimore, MD, 21287, USA). American Journal of Physiology, 265(5, Pt. 2), H1557--H1563 (English) 1993. CODEN: AJPHAP. ISSN: 0002-9513.

AB Large increases in intracranial pressure in fetal sheep result in more potent peripheral vasoconstriction and better maintenance of cerebral O<sub>2</sub> consumption (CMRO<sub>2</sub>) than in postnatal sheep. The fetus is exposed to a lower PO<sub>2</sub>. The authors tested the hypothesis that low PO<sub>2</sub> in postnatal lambs potentiates peripheral vasoconstriction and better maintains cerebral perfusion pressure and CMRO<sub>2</sub>. Pentobarbital-anesthetized lambs, 2-7 days old, were ventilated with either room air (n = 7) or a low O<sub>2</sub> mixture to reduce arterial O<sub>2</sub> saturation to 50% (n = 7). Elevation of intracranial pressure to within 3-5 mmHg of baseline mean arterial pressure for 30 min by ventricular fluid infusion initially caused a similar increase in arterial pressure in the normoxic (11 ± 3 (SE) mmHg) and hypoxic (14 ± 2 mmHg) groups. Plasma catecholamines increased more rapidly in the hypoxic group. However, plasma vasopressin levels were substantially elevated by hypoxia alone and failed to increase further with elevated intracranial pressure. Moreover, there was no significant difference between groups in the steady-state increase in arterial pressure, and microsphere-determined blood flow to intestines, kidney, skin, and muscle did not decrease in either group. Consequently, cerebral perfusion pressure, regional cerebral blood flow, and CMRO<sub>2</sub> were reduced similarly in both groups. Therefore, hypoxemia failed to potentiate the postnatal pressor response. Low PO<sub>2</sub> is unlikely to be the major mechanism for the potent Cushing response in the fetus.

L17 ANSWER 4 OF 8 EMBASE COPYRIGHT (c) 2008 Elsevier B.V. All rights reserved on STN

1993348201 EMBASE Effect of hypoxemia on the cardiovascular response to intracranial hypertension in postnatal lambs. Kearney M.L.; Backofen J.E.; Koehler R.C.; Jones Jr. M.D.; Traystman R.J.. R.C. Koehler, Anesthesiology/Crit. Care Med. Dept., Blalock 1404, Johns Hopkins Hospital, 600 N. Wolfe St., Baltimore, MD 21287-4961, United States. American Journal of Physiology - Heart and Circulatory Physiology Vol. 265, No. 5 34-5, pp. H1557-H1563 1993. ISSN: 0002-9513. CODEN: AJPPDI Pub. Country: United States. Language: English. Summary Language: English.

Entered STN: 931226. Last Updated on STN: 931226

AB Large increases in intracranial pressure in fetal sheep result in more potent peripheral vasoconstriction and better maintenance of cerebral O(2) consumption (CMR(O(2))) than in postnatal sheep. The fetus is exposed to a lower PO(2). We tested the hypothesis that low PO(2) in postnatal lambs potentiates peripheral vasoconstriction and better maintains cerebral perfusion pressure and CMR(O(2)). Pentobarbital-anesthetized lambs, 2-7 days old, were ventilated with either room air (n = 7) or a low O(2) mixture to reduce arterial O(2) saturation to 50% (n = 7). Elevation of intracranial pressure to within 3-5 mmHg of baseline mean arterial pressure for 30 min by ventricular fluid infusion initially caused a similar increase in arterial pressure in the normoxic ( $11 \pm 3$  (SE) mmHg) and hypoxic ( $14 \pm 2$  mmHg) groups. Plasma catecholamines increased more rapidly in the hypoxic group. However, plasma vasopressin levels were substantially elevated by hypoxia alone and failed to increase further with elevated intracranial pressure. Moreover, there was no significant difference between groups in the steady-state increase in arterial pressure, and microsphere-determined blood flow to intestines, kidney, skin, and muscle did not decrease in either group. Consequently, cerebral perfusion pressure, regional cerebral blood flow, and CMR(O(2)) were reduced similarly in both groups. Therefore, hypoxemia failed to potentiate the postnatal pressor response. Low PO(2) is unlikely to be the major mechanism for the potent Cushing response in the fetus.

L17 ANSWER 5 OF 8 EMBASE COPYRIGHT (c) 2008 Elsevier B.V. All rights reserved on STN

1993099552 EMBASE Transient hyponatremia after damage to the neurohypophyseal tracts. Ullmann M.C.; Hoffmann G.E.; Nelson P.B.; Robinson A.G.. Dr. A.G. Robinson, E-1140 Biomedical Science Tower, University of Pittsburgh, Pittsburgh, PA 15261, United States. Neuroendocrinology Vol. 56, No. 6, pp. 803-811 1992.

ISSN: 0028-3835. CODEN: NUNDAJ

Pub. Country: Switzerland. Language: English. Summary Language: English. Entered STN: 930516. Last Updated on STN: 930516

AB Section of the neurohypophyseal stalk classically produces a triphasic response: diabetes insipidus (1st phase), hyponatremia or normonatremia (2nd phase), and diabetes insipidus (3rd phase). Transient hyponatremia without diabetes insipidus has been reported after transphenoidal pituitary surgery. We report two additional cases of transient hyponatremia which occurred 6-8 days after pituitary surgery. We hypothesize that this outcome may be due to partial section or damage of the hypothalamic-neurohypophyseal tracts. The remaining intact vasopressin neurons function normally to protect against the diabetes insipidus of the first and third phase, but leak of vasopressin from the damaged tracts and posterior pituitary is sufficient to cause what can be described as an isolated second phase. To study this hypothesis in rats, partial damage to the hypothalamic-neurohypophyseal tracts was produced by radio-frequency lesions. The lesions did not affect anterior pituitary function. A variety of responses in posterior pituitary function occurred, including classic triphasic response in 2 rats and transient hyponatremia in 20 of 35 lesioned animals. The mean sodium nadir was  $128.7 \pm 1.5$  mEq/l in comparison to the sham-operated value of  $140.0 \pm 0.4$  mEq/l. Of the 20 rats exhibiting transient hyponatremia, 12 went on to develop diabetes insipidus, and 8 recovered. In the recovered group, the transient hyponatremia occurred 1-3 days after lesioning and returned to normal by day 7 which corresponds to the timing of the second phase of the triphasic response in rats. Hyponatremia was accompanied by vasopressin levels inappropriate for the plasma sodium level, inappropriately concentrated urine, water retention, and natriuresis. Animals that recovered from hyponatremia had sufficient vasopressin function to maintain normal plasma sodium with

normal levels of fluid intake and were able to tolerate 30 h of fluid deprivation with minimal dehydration and elevation of plasma sodium. However, in this group, the vasopressin release in response to hypertonic saline infusion was attenuated. At sacrifice immunohistochemistry of neurophysin was performed, and a portion of the hypothalamiconeurohypophyseal tract in the internal zone of the median eminence was found to be intact in all animals which recovered, i.e., there was partial section. Thus, in both the patients and in the animals, the transient hyponatremia had the characteristic etiology, timing, and duration of an isolated second phase of the triphasic response.

- L17 ANSWER 6 OF 8 MEDLINE on STN DUPLICATE 2  
 91064033. PubMed ID: 2147376. The role of arginine vasopressin in alcohol tolerance. Hoffman P L; Ishizawa H; Giri P R; Dave J R; Grant K A; Liu L I; Gulya K; Tabakoff B. (Division of Intramural Clinical and Biological Research, National Institute on Alcohol Abuse and Alcoholism, Bethesda, Maryland. ) Annals of medicine, (1990) Vol. 22, No. 4, pp. 269-74. Ref: 47. Journal code: 8906388. ISSN: 0785-3890. Pub. country: Finland. Language: English.
- AB Administration of the neuropeptide, arginine vasopressin, to animals that have acquired functional tolerance to ethanol will maintain such tolerance, even in the absence of further ethanol ingestion by the animals. In mice, this action of the peptide is mediated by central nervous system V1 receptors and requires intact brain noradrenergic systems. Autoradiographic studies have shown that some V1 receptors are localized presynaptically on catecholaminergic neuronal terminals in the mouse lateral septum, suggesting that vasopressin may act via modulation of catecholamine release. In addition, vasopressin has been found to increase mRNA levels for the proto-oncogene, c-fos, in septum and hippocampus, possibly by an action at postsynaptic receptors. Expression of c-fos, which has been hypothesized to play a role in central nervous system neuroadaptation, could transform short-term actions of vasopressin into long-term effects on ethanol tolerance. Studies with vasopressin antagonists indicate that the endogenous peptide influences tolerance, and therefore the effect of chronic ethanol ingestion on vasopressin synthesis and release was studied. In mice and rats, hypothalamic vasopressin mRNA is decreased by chronic ethanol exposure, although effects on plasma vasopressin levels differ in the two species. The effect of ethanol on extrahypothalamic vasopressin synthesis in brain is under investigation. The results suggest mechanisms by which vasopressin can produce long-term changes in central nervous system function, and provide evidence for a disturbance of vasopressin regulation during chronic ethanol ingestion.
- L17 ANSWER 7 OF 8 MEDLINE on STN DUPLICATE 3  
 90072400. PubMed ID: 2589491. Absent effect of plasma vasopressin on rat brain blood flow during hemorrhage. Nakai M; Yamane Y; Umeda Y; Inada M; Yamamoto J; Kawamura M. (Department of Cardiovascular Dynamics, National Cardiovascular Center Research Institute, Osaka, Japan. ) The American journal of physiology, (1989 Nov) Vol. 257, No. 5 Pt 2, pp. H1360-8. Journal code: 0370511. ISSN: 0002-9513. Pub. country: United States. Language: English.
- AB We investigated whether a reflex increase in plasma vasopressin level due to hemorrhagic hypotension affects brain blood flow. In 60 lightly anesthetized, artificially ventilated rats, the flow was determined with radiolabeled microspheres. We found excellent maintenance of blood flow throughout all brain regions during the hypotensive state (71 mmHg on average), and such maintenance of flow was not modulated at all by a supramaximal intravenous dose of the selective vasopressin V1-receptor antagonist [d(CH2)5 Tyr- (Me)AVP]. The latter finding also implies that the V1 antagonist failed to unmask the vasodilator type actions of V2 receptors on the maintenance of flow during

hemorrhagic hypotension. These were true also when the cervical sympathetic bundles were severed bilaterally. The plasma level of endogenous vasopressin was increased during hypotension, ranging from 118 to 973 pg/ml. Despite this increase, the brain blood flow was entirely independent of the plasma vasopressin level in all the brain regions studied. We conclude that the brain circulation of rats can maintain its blood flow during hemorrhagic hypotension without any apparent contribution from a concomitant reflex increase in plasma vasopressin. Despite our negative results for the brain blood flow, the possible segmental effects of circulating vasopressin on the brain arterial caliber remain to be clarified under conditions of hemorrhagic hypotension.

- L17 ANSWER 8 OF 8 EMBASE COPYRIGHT (c) 2008 Elsevier B.V. All rights reserved on STN
- 199009855 EMBASE Absent effect of plasma vasopressin on rat brain blood flow during hemorrhage. Nakai M.; Yamane Y.; Umeda Y.; Inada M.; Yamamoto J.; Kawamura M. M. Nakai, Dept. of Cardiovascular Dyn., Natl. Cardiov. Ctr. Res. Inst, Osaka 565, Japan. American Journal of Physiology - Heart and Circulatory Physiology Vol. 257, No. 5, pp. 26/5 1989. ISSN: 0002-9513. CODEN: AJPPDI  
Pub. Country: United States. Language: English. Summary Language: English. Entered STN: 911213. Last Updated on STN: 911213
- AB We investigated whether a reflex increase in plasma vasopressin level due to hemorrhagic hypotension affects brain blood flow. In 60 lightly anesthetized, artificially ventilated rats, the flow was determined with radiolabeled microspheres. We found excellent maintenance of blood flow throughout all brain regions during the hypotensive state (71 mmHg on average), and such maintenance of flow was not modulated at all by a supramaximal intravenous dose of the selective vasopressin V(1)-receptor antagonist [d(CH<sub>2</sub>)<sub>5</sub>Tyr-(Me)]AVP. The latter finding also implies that the V(1) antagonist failed to unmask the vasodilator type actions of V(2) receptors on the maintenance of flow during hemorrhagic hypotension. These were true also when the cervical sympathetic bundles were severed bilaterally. The plasma level of endogenous vasopressin was increased during hypotension, ranging from 118 to 973 pg/ml. Despite this increase, the brain blood flow was entirely independent of the plasma vasopressin level in all the brain regions studied. We conclude that the brain circulation of rats can maintain its blood flow during hemorrhagic hypotension without any apparent contribution from a concomitant reflex increase in plasma vasopressin. Despite our negative results for the brain blood flow, the possible segmental effects of circulating vasopressin on the brain arterial caliber remain to be clarified under conditions of hemorrhagic hypotension.

=> s blood vasopressin level  
L18 28 BLOOD VASOPRESSIN LEVEL

=> s l18 and unpredictable  
L19 0 L18 AND UNPREDICTABLE

=> s l18 and PTHrP  
L20 0 L18 AND PTHRP

=> s l18 nd anti-PTHrP  
MISSING OPERATOR L18 ND

The search profile that was entered contains terms or nested terms that are not separated by a logical operator.

=> s l18 and cancer

L21 0 L18 AND CANCER

=> dup remove 118

PROCESSING COMPLETED FOR L18

L22 18 DUP REMOVE L18 (10 DUPLICATES REMOVED)

=> s l22 and hypercalcemia

L23 0 L22 AND HYPERCALCEMIA

=> s l22 and pd<20000720

2 FILES SEARCHED...

L24 17 L22 AND PD<20000720

=> d l24 1-17 cbib abs

L24 ANSWER 1 OF 17 MEDLINE on STN

1998232569. PubMed ID: 9564057. Endometrial Na<sup>+</sup>, K<sup>+</sup>-ATPase pump function and vasopressin levels during hysteroscopic surgery in patients pretreated with GnRH agonist. Taskin O; Buhur A; Birincioglu M; Burak F; Atmaca R; Yilmaz I; Wheeler J M. (Department of Obstetrics and Gynecology, Inonu University Medical School, Malatya, Turkey. ) The Journal of the American Association of Gynecologic Laparoscopists, (1998 May) Vol. 5, No. 2, pp. 119-24. Journal code: 9417443. ISSN: 1074-3804. Pub. country: United States. Language: English.

AB STUDY OBJECTIVE: To investigate the effects of gonadotropin-releasing hormone (GnRH) analog pretreatment on endometrial Na<sup>+</sup>, K<sup>+</sup>-adenosine triphosphatase (ATPase) pump function and peripheral blood vasopressin levels, and their role in fluid absorption and mechanisms of hyponatremia in patients undergoing hysteroscopic endometrial ablation. DESIGN: Prospective, randomized, placebo-controlled study (Canadian Task Force classification I). SETTING: University-affiliated hospital. PATIENTS: Seventeen women with dysfunctional uterine bleeding. INTERVENTION: Nine women received a GnRH analog and eight received saline approximately 6 to 8 weeks before hysteroscopic ablation by electrosurgery. MEASUREMENTS and MAIN RESULTS: Both before randomization and immediately before surgery, endometrial biopsy samples were obtained and numbered consecutively without patient identification. Operative hysteroscopy was performed with glycine 1.5% mixed with 2% alcohol. The amount of irrigant and irrigant deficit; blood levels of albumin and ethanol; hematocrit and hemoglobin; changes in sodium levels; and central venous pressure were compared. The Na<sup>+</sup>, K<sup>+</sup>-ATPase pump activity was significantly increased in the GnRH analog group compared with the saline group and correlated with decreased estradiol levels (0.4 +/- 0.08 vs 0.26 +/- 0.06 micro mol/min/ml). Vasopressin levels were significantly lower in the GnRH group (3.2 +/- 0.9 vs 7.6 +/- 1.7 micro mol/L). Mean volume of irrigant used and operating time were similar in both groups. Volume deficit, decrease in protein, and hematocrit were less in GnRH than in the saline group. Blood ethanol levels, decrease in sodium, and irrigant deficit were significantly lower in GnRH group. CONCLUSION: Pretreatment with GnRH analogs may prevent the adverse effects of estradiol on endometrial Na<sup>+</sup>, K<sup>+</sup>-ATPase and creates a protective mechanism against iatrogenic hyponatremia, which is more critical in women than men in case of absorption of irrigating fluid. Moreover, created hypoestrogenism may enhance Na<sup>+</sup>, K<sup>+</sup>-ATPase activity in brain as well as endometrium, thus decreasing women's susceptibility to hyponatremic complications and brain damage. Suppressed vasopressin levels may be protective against fluid absorption in GnRH analog-treated patients.

L24 ANSWER 2 OF 17 MEDLINE on STN

96144056. PubMed ID: 8587172. Acute myelogenous leukemia with diabetes insipidus without desmopressin administration by anti-leukemic

chemotherapy. Ino Y; Tsurumi H; Yamada T; Murakami N; Moriwaki H; Muto Y. (First Department of Internal Medicine, Gifu University School of Medicine. ) [Rinsho ketsueki] The Japanese journal of clinical hematology, (1995 Dec) Vol. 36, No. 12, pp. 1359-64. Journal code: 2984782R. ISSN: 0485-1439. Pub. country: Japan. Language: Japanese.

- AB We report a case of AML with diabetes insipidus (DI). A 68-year-old female was admitted to our hospital because of fever and leukocytosis. The WBC was 197,000/microliter with 98% blasts positive for myeloperoxidase, CD33, CD34 and HLA-DR. While, on admission, urine volume was more than 6 liters daily. Blood vasopressin level was 0.3 microgram/ml. The patient was diagnosed as having AML with DI. By chemotherapy consisting of BHAC, DNR, 6-MP and PSL and intrathecal administration of AraC, MTX and PSL, and nasal drip of DDAVP, complete remission was attained and the urine volume was reduced to normal. Finally DDAVP became unnecessary. Although the exact cause of DI cannot be ascertained, rapid increase of leukemic blasts and leukostasis in small vessels might be associated with hypothalamus-pituitary system damage. Reportedly, DI is a rare complication of leukemia and administration of DDAVP could be halted in only two patients with leukemia and DI.

L24 ANSWER 3 OF 17 MEDLINE on STN

92177637. PubMed ID: 1795462. [Endogenous vasopressin and fibrinolysis in patients with angina pectoris]. Endogennyi vazopressin i fibrinoliz u bol'nykh so stenokardiei. Averkov O V; Zateishchikov D A; Gratsianskii N A; Dobrovol'skii A B; Panchenko E P; Masenko V P. Kardiologiya, (1991 Aug) Vol. 31, No. 8, pp. 11-4. Journal code: 0376351. ISSN: 0022-9040. Pub. country: USSR. Language: Russian.

- AB A relationship was examined between blood vasopressin levels and the fibrinolytic system in 35 patients with angina pectoris (16 with vasospastic angina (VA) and 19 with exercise-induced angina) who had undergone vein occlusion testing. There was a positive correlation between the post-testing vasopressin levels and the activity of tissue plasminogen activator inhibitor (TPAI) ( $r = 0.54$ ) which was more high in patients with VA ( $r = 0.61$ ). Only did the patients with VA show a direct relationship between the vasopressin concentrations and the activity of tissue plasminogen activator (TPA) ( $r = 0.63$ ), the concentration of fibrinogen-fibrin degradation products (FFDP) ( $r = 0.88$ ). Thirteen patients having higher vasopressin levels (over 3.4 ng/ml) displayed a greater TPAI activity than did the patients with vasopressin levels of at least 3.4 ng/ml ( $26.2 \pm 4.9$  and  $15.0 \pm 1.42$  IU/ml, respectively;  $p$  less than 0.05). There was a direct relationship between the vasopressin levels and the activity of TPA ( $r = 0.65$ ), the concentration of FFDP ( $r = 0.78$ ) in patients having a vasopressin level of above 3.4 ng/ml. The findings are in agreement with the concept that endogenous vasopressin is involved in the regulation of the blood fibrinolytic system.

L24 ANSWER 4 OF 17 MEDLINE on STN

91023165. PubMed ID: 2145777. Role of vasopressin in renal vascular changes with hypoxemia and hypercapnic acidosis in conscious dogs. Rose C E Jr; Ragsdale N V; Carey R M. (Department of Internal Medicine, University of Virginia School of Medicine, Charlottesville 22908. ) The American journal of physiology, (1990 Oct) Vol. 259, No. 4 Pt 2, pp. R690-702. Journal code: 0370511. ISSN: 0002-9513. Report No.: NASA-91023165. Pub. country: United States. Language: English.

- AB To evaluate the role of vasopressin in the renal changes during combined acute hypoxemia and acute hypercapnic acidosis, eight conscious female mongrel dogs prepared with controlled sodium intake at 80 meq/24 h for 4 days were studied in one of the following six protocols: acute hypoxemia (80 min, arterial PO2 34  $\pm$  1 mmHg) followed by combined acute hypoxemia and hypercapnic acidosis (40 min, arterial PO2 35  $\pm$  1 mmHg, arterial



PCO<sub>2</sub> 58 +/- 1 mmHg, pH = 7.20 +/- 0.01) during 1) intrarenal vehicle at 0.5 ml/min (N = 8); or 2) intrarenal infusion of vasopressin V1-receptor antagonist [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP at 5 ng.kg<sup>-1</sup>.min<sup>-1</sup> (N = 5); and with normal gas exchange during 3) intrarenal vasopressin at 0.05 mU.kg<sup>-1</sup>.min<sup>-1</sup> (N = 8); 4) simultaneous infusion of intrarenal vasopressin and [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP, 5 ng.kg<sup>-1</sup>.min<sup>-1</sup> (N = 4); 5) intrarenal [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP, 5 ng.kg<sup>-1</sup>.min<sup>-1</sup> (N = 4); and 6) intrarenal vehicle at 0.5 ml/min (N = 7). Intrarenal infusion of a suppressor dose of vasopressin resulted in a transient decrease in glomerular filtration rate and effective renal plasma flow over the first 20 min of infusion, suggesting that vasopressin induced nonsustained vasoconstriction of the renal vasculature. Intrarenal administration of [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP failed to block the fall in glomerular filtration rate or effective renal plasma flow when renal arterial blood vasopressin levels were elevated by intrarenal administration of exogenous vasopressin or by elevated systemic arterial endogenous circulating vasopressin during combined acute hypoxemia and hypercapnic acidosis. These data suggest that vasopressin (V1-receptor stimulation) does not play an important role in the renal vasoconstriction during combined acute hypoxemia and hypercapnic acidosis in conscious dogs.

- L24 ANSWER 5 OF 17 MEDLINE on STN  
 71137380. PubMed ID: 5401935. [Changes in the blood vasopressin level following certain balneophysiotherapeutic methods]. Modifications de la teneur en vasopressine sanguine apres certaines methodes balneophysiotherapeutiques. Modval M; Constantinescu J; Benetato V; Ionescu-Calinescu C. Revue roumaine de physiologie, (1969) Vol. 6, No. 4, pp. 285-91. Journal code: 7510990. ISSN: 0035-399X. Pub. country: Romania. Language: French.
- L24 ANSWER 6 OF 17 EMBASE COPYRIGHT (c) 2008 Elsevier B.V. All rights reserved on STN  
 1990358250 EMBASE Role of vasopressin in renal vascular changes with hypoxemia and hypercapnic acidosis in conscious dogs. Rose Jr. C.E.; Ragsdale N.V.; Carey R.M.. C.E. Rose Jr., Division of Pulmonary Medicine, Box 225, University of Virginia, Health Sciences Center, Charlottesville, VA 22908, United States. American Journal of Physiology - Regulatory Integrative and Comparative Physiology Vol. 259, No. 4 28-4, pp. R690-R702 1990. ISSN: 0002-9513. CODEN: AJPRDO Pub. Country: United States. Language: English. Summary Language: English. Entered STN: 911213. Last Updated on STN: 911213
- AB To evaluate the role of vasopressin in the renal changes during combined acute hypoxemia and acute hypercapnic acidosis, eight conscious female mongrel dogs prepared with controlled sodium intake at 80 meq/24 h for 4 days were studied in one of the following six protocols: acute hypoxemia (80 min, arterial PO<sub>2</sub>(2) 34 ± 1 mmHg) followed by combined acute hypoxemia and hypercapnic acidosis (40 min, arterial PO<sub>2</sub>(2) 35 ± 1 mmHg, arterial PCO<sub>2</sub>(2) 58 ± 1 mmHg, pH = 7.20 ± 0.01) during 1) intrarenal vehicle at 0.5 ml/min (N = 8); or 2) intrarenal infusion of vasopressin V(1)-receptor antagonist [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP at 5 ng.ovrhdot.kg(-1).ovrhdot.min(-1) (N = 5); and with normal gas exchange during 3) intrarenal vasopressin at 0.05 mU.ovrhdot.kg(-1).ovrhdot.min(-1) (N = 8); 4) simultaneous infusion of intrarenal vasopressin and [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP, 5 ng.ovrhdot.kg(-1).ovrhdot.min(-1) (N = 4); 5) intrarenal [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP, 5 ng.ovrhdot.kg(-1).ovrhdot.min(-1) (N = 4); and 6) intrarenal vehicle at 0.5 ml/min (N = 7). Intrarenal infusion of a suppressor dose of vasopressin resulted in a transient decrease in glomerular filtration rate and effective renal plasma flow over the first 20 min of infusion, suggesting that vasopressin induced nonsustained vasoconstriction of the renal vasculature. Intrarenal

administration of [d(CH<sub>2</sub>)<sub>5</sub>Tyr(Me)]AVP failed to block the fall in glomerular filtration rate or effective renal plasma flow when renal arterial blood vasopressin levels were elevated by intrarenal administration of exogenous vasopressin or by elevated systemic arterial endogenous circulating vasopressin during combined acute hypoxemia and hypercapnic acidosis. These data suggest that vasopressin (V(1)-receptor stimulation) does not play an important role in the renal vasoconstriction during combined acute hypoxemia and hypercapnic acidosis in conscious dogs.

- L24 ANSWER 7 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN 1978:245336 Document No.: PREV197866057833; BA66:57833. DEVICE FOR THE BIOASSAY OF VASOPRESSIN LEVEL IN BLOOD. HOLOVCHENKO S F [Reprint author]; MASHEK V O; NESTEROVS'KYI M V. LAB PHYSIOL, INST GERONTOL, ACAD MED SCI USSR, KIEV, USSR. Fiziologichnyi Zhurnal (Kiev), (1977) Vol. 23, No. 5, pp. 707-708. CODEN: FZUKAM. ISSN: 0015-3311. Language: UKRAINIAN.
- AB Bioassay of human blood vasopressin levels on rats is based on its diuretic effect. A 5-channel device for registering diuresis continuously for 6-10 h with an accuracy of 0.02 ml is described. The device can also be used for the graphic registration of drops of other electroconductive fluids.
- L24 ANSWER 8 OF 17 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN 1978:171161 Document No.: PREV197865058161; BA65:58161. STUDIES ON THE ROLE OF HIGH PRESSURE BARO RECEPTORS IN VASOPRESSIN SECRETION EFFECT OF OCCLUSION OF COMMON CAROTID AND VERTEBRAL ARTERIES ON BLOOD VASOPRESSIN LEVEL. MATSUZAKI M [Reprint author]. SECOND DEP SURG, NAGOYA UNIV SCH MED, NAGOYA, AICHI, JPN. Folia Endocrinologica Japonica, (1977) Vol. 53, No. 8, pp. 982-989. CODEN: NNGZAZ. ISSN: 0029-0661. Language: JAPANESE.
- AB The role of baroreceptors, in common carotid and vertebral arteries and arteries in the thoracic cavity, in vasopressin [ADH] secretion was investigated. Effects of bilateral occlusion of the common carotid and vertebral arteries on blood ADH levels and mean arterial pressure [MAP] were studied in common carotid arterial plexus-denervated dogs, cervically vagotomized dogs and intact dogs. Blood ADH titers were determined by bioassay before and 5 min after the occlusion of the arteries and were compared with the changes of MAP. Blood ADH titers and MAP were elevated by the occlusion of the common carotid arteries in intact and vagotomized dogs, while they were not significantly affected in denervated dogs. Elevation of blood ADH titers was more pronounced in vagotomized dogs than in intact dogs. Blood ADH titers and MAP were elevated by the occlusion of vertebral arteries in all dogs. The elevation of blood ADH titers in denervated dogs was more pronounced than in intact dogs, but less than in vagotomized dogs. The effects of common carotid artery occlusion on blood ADH titers and MAP were more pronounced than those of the vertebral artery occlusion. Baroreceptors involved in vasopressin secretion are present in vertebral arteries and the intrathoracic baroreceptors are dominant in controlling vasopressin secretion, while those in common carotid arteries are of secondary importance and those in vertebral arteries are less important.
- L24 ANSWER 9 OF 17 SCISEARCH COPYRIGHT (c) 2008 The Thomson Corporation on STN 1992:677254 The Genuine Article (R) Number: JX717. EFFECT OF PARATHYROID-HORMONE ON CA-45(2+) ACCUMULATION NEUROSECRETORY-CELLS AND ON BLOOD VASOPRESSIN LEVELS AFTER PARATHYROIDECTOMY AND INJECTION OF PARATHYROID EXTRACT. KHUVERDYAN D N (Reprint); ASRATYAN A A. YEREVAN PHYS INST, DEPT NORMAL PHYSIOL, YEREVAN, ARMENIA, USSR (Reprint). BULLETIN OF EXPERIMENTAL BIOLOGY AND MEDICINE (MAR 1992) Vol. 113, No. 3, pp. 289-291. ISSN: 0007-4888.

Publisher: PLENUM PUBL CORP, CONSULTANTS BUREAU 233 SPRING ST, NEW YORK, NY 10013. Language: English.

L24 ANSWER 10 OF 17 SCISEARCH COPYRIGHT (c) 2008 The Thomson Corporation on STN

1986:525566 The Genuine Article (R) Number: D9177. BLOOD VASOPRESSIN LEVELS IN RELATION TO OTHER HORMONES IN CORONARY PATIENTS. DUDAIEV V A (Reprint); GORIN V V; BORODKIN V V; DYUKOV I V; NECHAEVA N I. NI PIROGOV MED INST, FAC MED, DEPT INTERNAL DIS 1, MOSCOW, USSR (Reprint). KARDIOLOGIYA (JUL 1986) Vol. 26, No. 7, pp. 98-101. ISSN: 0022-9040. Publisher: IZD VO MEDITSINA, PETROVERIGSKII PER 6-8, K-142 MOSCOW, RUSSIA. Language: Russian.

L24 ANSWER 11 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

1997:297524 Document No. 126:272593 Calcium-parathyroid hormone system in the functional activity of the hypothalamic-neurohypophyseal complex. Khudaverdyan, D. N.; Asratyan, A. A. (Erevan. Gos. Med. Univ., Yerevan, Armenia). Byulleten Eksperimental'noi Biologii i Meditsiny, 122(11), 484-486 (Russian) 1996. CODEN: BEBMAE. ISSN: 0365-9615. Publisher: Meditsina.

AB The effect of single and multiple i.m. administration of parathormone on protein synthesis in neurosecretory cells of the supraoptic nucleus and blood vasopressin content was investigated in rats. Parathormone administration increased RNA expression in supraoptic nucleus cells and blood vasopressin levels, especially on single administration.

L24 ANSWER 12 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

1992:34845 Document No. 116:34845 Hormonal indexes of the reactivity of central parts of hypothalamo-hypophyseal-adrenocortical system in rats during postnatal life. Danilova, O. A.; Chernigovskaya, E. V.; Chetverukhin, V. K. (Inst. Evol. Physiol. Biochem., Leningrad, USSR). Zhurnal Evolyutsionnoi Biokhimii i Fiziologii, 27(3), 308-13 (Russian) 1991. CODEN: ZEBFAJ. ISSN: 0044-4529.

AB ACTH and vasopressin were determined in the blood and pituitary glands of adult male rats as well as in 1-, 3-, 5-, 7-, and 20-day-old rat pups and in the pups at 30 min after surgical stress (cutting skin on the back). The ACTH level in the blood at 1 day of age was 25 pg/mL increasing to a maximum of 279 pg/mL in the adult. Pituitary ACTH levels reached a peak of 17,500 pg/100 g at 20 days of age. Blood vasopressin showed peak levels of 13.0 and 14.2 pg/mL in 5-day-old and adult rats; pituitary levels generally decreased from birth to adult. Surgical stress decreased ACTH secretion in 1-5-day-old rats, increased it in 7-day-old animals, and decreased it in 20-day-old and adult animals. Pituitary levels of ACTH increased in 3-day-old animals and decreased in 7-day-old animals in response to stress. Blood vasopressin levels were decreased in 3-day-old rats, but were increased in 7-day-old rats in response to stress. In the pituitary, vasopressin levels were increased in 5- and 20-day-old rats. Vasopressin increased in the median eminence of 1-, 3-, and 20-day old rats in response to stress. Thus, the early period of development of the adrenal cortex-hypothalamus-pituitary system is characterized by a paradoxical reaction rather than by nonreactivity which may be associated with poor neurohormonal transport in the outer zone of the median eminence.

L24 ANSWER 13 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

1988:469548 Document No. 109:69548 The content of vasopressin in the rabbit's blood plasma after the action of ionizing radiation. Gzirishvili, N. A.; Kbuladze, G. I.; Kabzinadze, K. G. (I. S. Beritashvili Inst. Physiol., Tbilisi, USSR). Izvestiya Akademii Nauk Gruzinskoi SSR, Seriya Biologicheskaya, 14(2), 82-7 (Russian) 1988. CODEN: IGSBDO. ISSN: 0321-1665.

AB Whole body exposure of rabbits to x-irradiation (4-8 Gy) or local x-irradiation of the head (8 Gy) altered vasopressin levels in the blood. At 30 min after both whole body and local irradiation, blood vasopressin levels were decreased; this was followed, at 2 h, by a marked increase above initial levels. In rabbits with more severe irradiation damage (whole body and local exposure to 8 Gy x-irradiation) blood vasopressin levels were elevated between 5 and 30 days after exposure, whereas in animals exposed to whole body-irradiation at 4 Gy they were below normal between 10 and 30 days after exposure. These changes in vasopressin secretion following irradiation may be related to the regulation of metabolic processes in the development of adaptive, compensatory, and reparative processes during radiation disease.

L24 ANSWER 14 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

1985:572535 Document No. 103:172535 Original Reference No. 103:27583a,27586a Vasopressin effect on the cardiovascular system and electrical activity of the hypothalamus in rabbits of different age. Pugach, B. V. (Inst. Gerontol., Kiev, USSR). Fiziologicheskii Zhurnal (Kiev, 1978-1993), 31(4), 429-33 (Russian) 1985. CODEN: FIZHDO. ISSN: 0201-8489.

AB Variations in the hemodynamics and elec. activity of the hypothalamus evoked by a single i.v. injection of vasopressin [11000-17-2] (0.2 units/kg) into 10-12- and 48-60-mo-old rabbits were studied. Vasopressin increased arterial pressure and general peripheral resistance, decreased heart min. volume, caused bradycardia, and depressed the S-T segment of the electrocardiograph. The cardiovascular effects were usually more expressed in senescent than in mature animals. Vasopressin also altered (generally decreased) the elec. activity of the supraoptic and ventromedial nuclei of the hypothalamus, effects which were usually more expressed in mature than in senescent animals. The results were related to the increase in blood vasopressin levels observed in senescence.

L24 ANSWER 15 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

1970:19760 Document No. 72:19760 Original Reference No. 72:3589a,3592a Vasopressin content in the hypothalamus, hypophysis, and blood plasma in the guinea pig. Guzek, Jan W. (Sch. Med., Lodz, Pol.). Annals of the Medical Section of the Polish Academy of Sciences, 13(2), 175-205 (English) 1969. CODEN: ALMPBF. ISSN: 0048-4733.

AB The vasopressin content in the hypothalamus, hypophysis and blood plasma was determined in hydrated and dehydrated guinea pigs under various conditions of adrenergic transmission. In the hydrated guinea pigs, plasma vasopressin level drops below the threshold sensitivity of the assay. After 5 days of dehydration, a marked decrease in the vasopressin content in the hypothalamus accompanied by a distinct rise in the blood level was observed. A single dose of reserpine and amphetamine, resp., causes, in hydrated guinea pigs, a marked diminution in the vasopressin content in the hypothalamus and an increase in the blood vasopressin level. Reserpine and amphetamine administered to guinea pigs dehydrated for 5 days result, in an increase of the vasopressin release from the hypothalamo-hypophyseal system as compared with animals simply dehydrated. It is supposed that the increased vasopressin release following reserpine administration might be due to the inhibition of adrenergic inhibitory neurons. Similarly, increased vasopressin liberation which follows treatment with amphetamine, might be due to the activation of adrenergic excitatory neurons.

L24 ANSWER 16 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

1969:409946 Document No. 71:9946 Original Reference No. 71:1819a,1822a Release of vasopressin in response to hemorrhage and its role in the mechanism of blood pressure regulation. Rocha e Silva, Mauricio; Rosenberg, Manuela (Med. Sch., Sao Paulo Univ., Sao Paulo, Brazil).

Journal of Physiology (Cambridge, United Kingdom), 202, 535-57 (English) 1969. CODEN: JPHYA7. ISSN: 0022-3751.

- AB The release of vasopressin in response to hemorrhage and the effects of vasopressin infusions on blood pressure and heart rate were studied in dogs. Decrease in diastolic blood pressure of 21-30 mm. elicited increases in blood levels of vasopressin from a control value of 12.8 to 64.8 and 28.5 microunits/ml. in 5- and 30-min. hemorrhage blood samples, resp. Re-transfusion of blood restored vasopressin to control levels. Infusions of vasopressin in amts. secreted in response to hemorrhage evoked vasopressor responses when blood-pressure regulating reflexes were suppressed in reserpinized and atropinized dogs. The role of the secretion of endogenous vasopressin by the pituitary gland in the regulation of blood pressure was also studied in hypophysectomized and deafferented (bilaterally divided vagal and sinus nerves) dogs. Hypotensive responses were paralleled by decreases in blood vasopressin levels. Therefore, the release of vasopressin in response to stimuli from cardiovascular sensory receptors is involved in blood-pressure regulation.

L24 ANSWER 17 OF 17 CAPLUS COPYRIGHT 2008 ACS on STN

1965:473730 Document No. 63:73730 Original Reference No. 63:13648f-g The effects of calcium on protein-binding and metabolism of arginine vasopressin in rats. Smith, M. W.; Thorn, N. A. (Univ. Copenhagen). Journal of Endocrinology, 32(2), 141-51 (Unavailable) 1965. CODEN: JOENAK. ISSN: 0022-0795.

- AB cf. CA 63, 8923b. In rats made hypercalcemic by intravenous CaCl<sub>2</sub> injection, then given vasopressin intravenously, blood vasopressin levels fell more slowly than in normocalcemic controls. In the 30 min. following injection, average urinary excretion by controls was equivalent to 7% of the vasopressin given, and that by hypercalcemic rats was 24%. In controls, injected vasopressin was distributed in a volume equal to blood volume, but in hypercalcemic rats, the distribution volume was about 3 times as large. Antidiuresis from injection of large amts. of vasopressin into hydrated rats was little affected by blood Ca changes. Intravenous CaCl<sub>2</sub> given hydrated rats caused temporary diuresis. Expts. with Sephadex G-25 in vitro showed that both ox neurophysin and rat serum protein bind vasopressin, and that Ca interferes.

=> s (ogalta e?/au or onuma e?/au or tsunenari t?/au or saito h?/au or azuma y?/au)  
L25 35483 (OGALTA E7/AU OR ONUMA E7/AU OR TSUNENARI T7/AU OR SAITO H7/AU  
OR AZUMA Y7/AU)

=> s l25 and humanized anti-PTHrP  
L26 14 L25 AND HUMANIZED ANTI-PTHRP

=> dup remove l26  
PROCESSING COMPLETED FOR L26  
L27 4 DUP REMOVE L26 (10 DUPLICATES REMOVED)

=> d l27 1-4 cbib abs

L27 ANSWER 1 OF 4 MEDLINE on STN DUPLICATE 1  
2005628438. PubMed ID: 16309168. Humanized monoclonal antibody against parathyroid hormone-related protein suppresses osteolytic bone metastasis of human breast cancer cells derived from MDA-MB-231. Saito Hidemi ; Tsunenari Toshiaki; Onuma Etsuro; Sato Koh; Ogata Etsuro; Yamada-Okabe Hisafumi. (Pharmaceutical Research Department III, Kamakura Research Laboratories, Chugai Pharmaceutical Co., Ltd., Kamakura, Kanagawa 247-8530, Japan. ) Anticancer research, (2005 Nov-Dec) Vol. 25, No. 6B, pp. 3817-23. Journal code: 8102988. ISSN: 0250-7005. Pub.

country: Greece. Language: English.

- AB BACKGROUND: Parathyroid hormone-related protein (PTHrP) has been implicated in bone metastasis. However, the effects on bone metastatic activity of blocking the PTHrP function have not been tested in the clinic. Here, the effects of a humanized anti-PTHrP monoclonal antibody (mAb) on bone metastasis in a human xenograft model are shown. MATERIALS AND METHODS: Subline MDA-5a, with high bone metastatic activity, was established from the human breast cancer cell line MDA-MB-231. Mice were injected with MDA-5a and an anti-PTHrP monoclonal antibody (mAb) raised against human PTHrP (1-34); bone metastasis was evaluated by X-ray photography. RESULTS: MDA-5a produced elevated levels of PTHrP, Interleukin 8 (IL-8), IL-6 and matrix metalloproteinase 1 (MMP-1) and frequently metastasized to the bone. Administration of the humanized anti-PTHrP mAb significantly suppressed osteolytic bone metastasis of MDA-5a and caused osteogenesis at the sites of metastasis. CONCLUSION: The humanized anti-PTHrP mAb was effective against bone metastasis by inducing osteogenesis and, therefore, will provide a new treatment option for bone metastasis in breast cancer.

- L27 ANSWER 2 OF 4 MEDLINE on STN DUPLICATE 2  
2004546213. PubMed ID: 15517871. Generation of a humanized monoclonal antibody against human parathyroid hormone-related protein and its efficacy against humoral hypercalcemia of malignancy. Onuma Etsuro ; Sato Koh; Saito Hidemi; Tsunenari Toshiaki; Ishii Kimie; Esaki Keiko; Yabuta Naohiro; Wakahara Yuji; Yamada-Okabe Hisafumi; Ogata Etsuro. (Chugai Research Laboratories, Chugai Pharmaceutical Co. Ltd., 200 Kajiwara, Kamakura, Kanagawa, Japan. ) Anticancer research, (2004 Sep-Oct) Vol. 24, No. 5A, pp. 2665-73. Journal code: 8102988. ISSN: 0250-7005. Pub. country: Greece. Language: English.

- AB A humanized monoclonal antibody against parathyroid hormone-related protein (PTHrP) was generated from the mouse monoclonal antibody raised against the peptide corresponding to the N-terminal 34 amino acids of the human PTHrP [(PTHrP(1-34))]. The humanized antibody interacted with the PTHrP(1-34) with a kD value of  $1.90 \times 10^{(-10)}$  M, and the epitope resides between the amino acids 20 and 30 of the PTHrP. PTHrP(1-34) significantly increased the intracellular cAMP levels in the rat osteosarcoma cells that expressed PTHrP, and the 5 microg/mL or higher concentrations of the humanized antibody almost completely blocked the PTHrP-induced cAMP production even in the presence of 2 microg/mL PTHrP(1-34), demonstrating its ability to fully neutralize PTHrP function. There was no significant difference in the potency of the mouse, chimera, or the humanized antibodies to suppress the PTHrP-induced increase in the intracellular cAMP in ROS cells. Furthermore, at the same doses, the administration of the chimera or the humanized antibody was equally effective in reducing the blood ionized calcium levels of hypercalcemic mice bearing the PAN-7-JCK human pancreatic cancer xenograft or the LC-6-JCK human lung cancer xenograft that secreted PTHrP. Thus, humanized anti-PTHrP may be useful for the treatment of the humoral hypercalcemia of malignancy in humans.

- L27 ANSWER 3 OF 4 MEDLINE on STN DUPLICATE 3  
2003534792. PubMed ID: 14613038. Treatment of malignancy-associated hypercalcemia and cachexia with humanized anti-parathyroid hormone-related protein antibody. Sato Koh; Onuma Etsuro; Yocum Richard C; Ogata Etsuro. (Department of International Coordination, Chugai Pharmaceutical Co., Ltd, Skizuuoka, Japan. ) Seminars in oncology, (2003 Oct) Vol. 30, No. 5 Suppl 16, pp. 167-73. Ref: 11. Journal code: 0420432. ISSN: 0093-7754. Pub. country: United States. Language: English.

- AB Parathyroid hormone-related protein (PTHrP) plays a central role in humoral hypercalcemia of malignancy (HHM), which is one of the most frequent paraneoplastic syndromes. PTHrP produced by the tumor acts

through a common PTH/PTHrP receptor to promote bone resorption, inhibit calcium excretion from the kidney, and induce hypercalcemia. Patients with HHM often develop cachexia associated with typical symptoms such as anorexia, malaise, nausea, constipation, polyuria, polydipsia, and confusion. The etiology of the cachexia is not fully understood but is thought to be caused by hypercalcemia and various cytokines such as interleukin-6, tumor necrosis factor- $\alpha$ , leukemia inhibitory factor, and others. In this study, we investigated the role of PTHrP in hypercalcemia and cachexia in HHM by using humanized anti-PTHrP antibody. A mouse monoclonal antibody that binds to PTHrP amino acid sequence 1-34 and inhibits PTHrP function has been humanized to create a specific and potent agent for the treatment of patients with HHM. The mouse monoclonal antibody has been shown to have antihypercalcemic activity against nude mice bearing human tumors. Because a mouse antibody is highly immunogenic in human patients, the complementarity-determining regions from the mouse antibody were grafted into a human antibody. The resulting humanized antibody specifically recognizes PTHrP(1-34) and neutralizes PTHrP functions in vitro and in vivo. The humanized anti-PTHrP antibody was administered intravenously to HHM model animals bearing tumors such as LC-6 human lung carcinoma. These animals showed symptoms similar to those of patients with HHM (eg, hypercalcemia and cachexia). The humanized anti-PTHrP antibody-treated animals responded with normalization of blood ionized calcium level through an improvement of bone metabolism and calcium excretion. Moreover, the treated animals also showed an improvement in body weight, ultramotility, metabolic alkalosis, food consumption, water intake, serum phosphorus, and renal function. Consequently, the humanized antibody-treated animals experienced complete resolution of hypercalcemia and cachexia. These results suggest that the humanized antibody would be an effective and beneficial agent for patients with HHM, and that PTHrP is a major pathogenetic factor of hypercalcemia and cachexia in patients with HHM.

L27 ANSWER 4 OF 4 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN  
2000:244979 Document No.: PREV200000244979. The possibility of utilizing  
humanized anti-PTHrP antibody as an  
anti-HHM/cachexia agent. Onuma, Etsuro [Reprint author];  
Saito, H.; Azuma, Y.; Shimizu, N.; Tsunenari, T.  
; Sato, K.; Ogata, E.. Chugai Pharmaceutical, Shizuoka, Japan. Proceedings  
of the American Association for Cancer Research Annual Meeting, (March,  
2000) No. 41, pp. 287. print.  
Meeting Info.: 91st Annual Meeting of the American Association for Cancer  
Research. San Francisco, California, USA. April 01-05, 2000.  
ISSN: 0197-016X. Language: English.

=> s 125 and "FERM BP-5631"  
L28 0 L25 AND "FERM BP-5631"

=> s "FERM BP-5631"  
L29 0 "FERM BP-5631"

=> s 125 and hybridoma  
L30 79 L25 AND HYBRIDOMA

=> s 130 and "23-57-137-1"  
L31 1 L30 AND "23-57-137-1"

=> d 131 cbib abs

L31 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN  
2002:888597 Document No. 138:3671 Angiogenesis inhibitors that block binding

of PTH-related peptide to its receptor for use as antitumor agents.  
 Saito, Hidemi; Tsunenari, Toshiaki; Onuma,  
 Etsuro; Kato, Atsuhiko; Suzuki, Masami (Chugai Seiyaku Kabushiki  
 Kaisha, Japan). PCT Int. Appl. WO 2002092133 A1 20021121, 110 pp.  
 DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ,  
 CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE,  
 GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
 LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO,  
 RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,  
 YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ,  
 CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC,  
 ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (Japanese). CODEN: PIXXD2.  
 APPLICATION: WO 2002-JP4586 20020510. PRIORITY: JP 2001-140659 20010510.

AB It is found out that angiogenesis can be inhibited by a substance which  
 inhibits the binding of a parathyroid hormone-associated peptide (e.g. PTHrP)  
 to its receptor. The angiogenesis inhibitors can be anti-PTHrP  
 antibodies, antibody fragments, humanized or chimeric antibodies, PTH  
 receptor antagonists, or antisense oligonucleotides specific to PTHrP.  
 These modified anti-PTHrP antibodies and PTH receptor antagonists are  
 useful as antitumor agents and bone metastasis inhibitors.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	197.56	197.77
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-10.40	-10.40
STN INTERNATIONAL LOGOFF AT 09:34:49 ON 24 JAN 2008		

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSSPTA1644PNH

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1

Web Page for STN Seminar Schedule - N. America



NEWS 2 AUG 15 CAOLD to be discontinued on December 31, 2008  
 NEWS 3 OCT 07 EPFULL enhanced with full implementation of EPC2000  
 NEWS 4 OCT 07 Multiple databases enhanced for more flexible patent  
 number searching  
 NEWS 5 OCT 22 Current-awareness alert (SDI) setup and editing  
 enhanced  
 NEWS 6 OCT 22 WPIDS, WPINDEX, and WPIX enhanced with Canadian PCT  
 Applications  
 NEWS 7 OCT 24 CHEMLIST enhanced with intermediate list of  
 pre-registered REACH substances  
 NEWS 8 NOV 21 CAS patent coverage to include exemplified prophetic  
 substances identified in English-, French-, German-,  
 and Japanese-language basic patents from 2004-present  
 NEWS 9 NOV 26 MARPAT enhanced with F\$ORT command  
 NEWS 10 NOV 26 MEDLINE year-end processing temporarily halts  
 availability of new fully-indexed citations  
 NEWS 11 NOV 26 CHEMSAFE now available on STN Easy  
 NEWS 12 NOV 26 Two new SET commands increase convenience of STN  
 searching  
 NEWS 13 DEC 01 ChemPort single article sales feature unavailable  
 NEWS 14 DEC 12 GBFULL now offers single source for full-text  
 coverage of complete UK patent families

NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,  
 AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
 NEWS LOGIN Welcome Banner and News Items  
 NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that  
 specific topic.

All use of STN is subject to the provisions of the STN Customer  
 agreement. Please note that this agreement limits use to scientific  
 research. Use for software development or design or implementation  
 of commercial gateways or other similar uses is prohibited and may  
 result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:52:25 ON 16 DEC 2008

=> file medline embase biosis scisearch caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 12:52:50 ON 16 DEC 2008

FILE 'EMBASE' ENTERED AT 12:52:50 ON 16 DEC 2008  
 Copyright (c) 2008 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 12:52:50 ON 16 DEC 2008  
 Copyright (c) 2008 The Thomson Corporation

FILE 'SCISEARCH' ENTERED AT 12:52:50 ON 16 DEC 2008  
 Copyright (c) 2008 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 12:52:50 ON 16 DEC 2008  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

=> s blood vasopressin levels

L1 24 BLOOD VASOPRESSIN LEVELS

=> s l1 and parathyroid hormone

L2 2 L1 AND PARATHYROID HORMONE

=> dup remove l2

PROCESSING COMPLETED FOR L2

L3 2 DUP REMOVE L2 (0 DUPLICATES REMOVED)

=> d l3 1-2 cbib abs

L3 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2008 ACS on STN

1997:297524 Document No. 126:272593 Original Reference No. 126:52697a,52700a

Calcium-parathyroid hormone system in the functional activity of the hypothalamic-neurohypophyseal complex. Khudaverdyan, D. N.; Asratyan, A. A. (Erevan. Gos. Med. Univ., Yerevan, Armenia). Byulleten Eksperimental'noi Biologii i Meditsiny, 122(11), 484-486 (Russian) 1996. CODEN: BEBMAE. ISSN: 0365-9615. Publisher: Meditsina.

AB The effect of single and multiple i.m. administration of parathormone on protein synthesis in neurosecretory cells of the supraoptic nucleus and blood vasopressin content was investigated in rats. Parathormone administration increased RNA expression in supraoptic nucleus cells and blood vasopressin levels, especially on single administration.

L3 ANSWER 2 OF 2 SCISEARCH COPYRIGHT (c) 2008 The Thomson Corporation on STN

1992:677254 The Genuine Article (R) Number: JX717. EFFECT OF

PARATHYROID-HORMONE ON CA-45(2+) ACCUMULATION NEUROSECRETORY-CELLS AND ON BLOOD VASOPRESSIN LEVELS AFTER PARATHYROIDECTOMY AND INJECTION OF PARATHYROID EXTRACT. KHUDAVERDYAN D N (Reprint); ASRATYAN A A. YEREVAN PHYS INST, DEPT NORMAL PHYSIOL, YEREVAN, ARMENIA, USSR (Reprint). BULLETIN OF EXPERIMENTAL BIOLOGY AND MEDICINE (MAR 1992) Vol. 113, No. 3, pp. 289-291. ISSN: 0007-4888. Publisher: PLENUM PUBL CORP, CONSULTANTS BUREAU 233 SPRING ST, NEW YORK, NY 10013. Language: English.

=> s anti-PThrP

L4 261 ANTI-PTHRP

=> s l4 and increase

L5 66 L4 AND INCREASE

=> s l5 and blood vasopressin levels

L6 0 L5 AND BLOOD VASOPRESSIN LEVELS

=> s l4 and polyuria

L7 4 L4 AND POLYURIA

=> dup remove l7

PROCESSING COMPLETED FOR L7

L8 2 DUP REMOVE L7 (2 DUPLICATES REMOVED)

=> d l8 1-2 cbib abs

L8 ANSWER 1 OF 2 MEDLINE on STN

200605729. PubMed ID: 16444366. [Hypercalcemia of malignancy: clinical features, diagnosis and treatment]. A hipercalcemia nas malignidades:

aspectos clinicos, diagnosticos e terapeuticos. Farias Maria Lucia F de. (Faculdade de Medicina, Universidade Federal do Rio de Janeiro, RJ.. fleiuss@hucff.ufrj.br) . Arquivos brasileiros de endocrinologia e metabologia, (2005 Oct) Vol. 49, No. 5, pp. 816-24. Electronic Publication: 2006-01-23. Ref: 64. Journal code: 0403437. ISSN: 0004-2730. Pub. country: Brazil. Language: Portuguese.

- AB Hypercalcemia associated with malignancies is reported in up to 20 to 30% of patients with cancer during the course of the disease, and points to a poor prognosis. Symptoms related to the central nervous system, as progressive mental impairment, stupor and coma, predominate. Alterations in kidney function (water-concentrating defect leading to polyuria) and gastrointestinal tract (anorexia, nausea, vomiting) corroborate to dehydration and a further increase in serum calcium. Cancer-induced hypercalcemia may be classified as: 1) local osteolytic hypercalcemia (LOH), due to marked increase in osteoclastic bone resorption in areas surrounding the malignant cells within the marrow space; 2) humoral hypercalcemia of malignancy, caused by the secretion of parathyroid hormone-related protein (PTHrP) by the malignant tumor; 3) ectopic hyperparathyroidism; 4) 1,25(OH)<sub>2</sub> D-secreting tumors. Adequate control of hypercalcemia is necessary to give the patient time to respond to anti-cancer therapy. Volume expansion with saline will correct dehydration, improve glomerular filtration and increase urinary calcium excretion, which may be further stimulated by loop diuretics. Intravenous bisphosphonates are the most effective agents to control hypercalcemia, as they block osteoclastic osteolysis and also have antitumoral effects, decreasing bone metastases. New approaches to control the skeletal manifestations of malignancies are anti-PTHrP and anti-RANKL antibodies, osteoprotegerin, and also proteasome inhibitors in the case of multiple myeloma.

L8 ANSWER 2 OF 2 MEDLINE on STN DUPLICATE 1  
2003534792. PubMed ID: 14613038. Treatment of malignancy-associated hypercalcemia and cachexia with humanized anti-parathyroid hormone-related protein antibody. Sato Koh; Onuma Etsuro; Yocum Richard C; Ogata Etsuro. (Department of International Coordination, Chugai Pharmaceutical Co, Ltd, Skizuoka, Japan. ) Seminars in oncology, (2003 Oct) Vol. 30, No. 5 Suppl 16, pp. 167-73. Ref: 11. Journal code: 0420432. ISSN: 0093-7754. Pub. country: United States. Language: English.

- AB Parathyroid hormone-related protein (PTHrP) plays a central role in humoral hypercalcemia of malignancy (HHM), which is one of the most frequent paraneoplastic syndromes. PTHrP produced by the tumor acts through a common PTH/PTHrP receptor to promote bone resorption, inhibit calcium excretion from the kidney, and induce hypercalcemia. Patients with HHM often develop cachexia associated with typical symptoms such as anorexia, malaise, nausea, constipation, polyuria, polydipsia, and confusion. The etiology of the cachexia is not fully understood but is thought to be caused by hypercalcemia and various cytokines such as interleukin-6, tumor necrosis factor- $\alpha$ , leukemia inhibitory factor, and others. In this study, we investigated the role of PTHrP in hypercalcemia and cachexia in HHM by using humanized anti-PTHrP antibody. A mouse monoclonal antibody that binds to PTHrP amino acid sequence 1-34 and inhibits PTHrP function has been humanized to create a specific and potent agent for the treatment of patients with HHM. The mouse monoclonal antibody has been shown to have antihypercalcemic activity against nude mice bearing human tumors. Because a mouse antibody is highly immunogenic in human patients, the complementarity-determining regions from the mouse antibody were grafted into a human antibody. The resulting humanized antibody specifically recognizes PTHrP(1-34) and neutralizes PTHrP functions in vitro and in vivo. The humanized anti-PTHrP antibody was administered intravenously to HHM model animals bearing tumors such as LC-6 human lung carcinoma. These animals showed symptoms similar to those of patients with HHM (eg,

hypercalcemia and cachexia). The humanized anti-PTHrP antibody-treated animals responded with normalization of blood ionized calcium level through an improvement of bone metabolism and calcium excretion. Moreover, the treated animals also showed an improvement in body weight, ultramotivity, metabolic alkalosis, food consumption, water intake, serum phosphorus, and renal function. Consequently, the humanized antibody-treated animals experienced complete resolution of hypercalcemia and cachexia. These results suggest that the humanized antibody would be an effective and beneficial agent for patients with HHM, and that PTHrP is a major pathogenetic factor of hypercalcemia and cachexia in patients with HHM.

=> s l4 and dehydration  
L9 1 L4 AND DEHYDRATION

=> d l9 cbib abs

L9 ANSWER 1 OF 1 MEDLINE on STN  
2006057279. PubMed ID: 16444366. [Hypercalcemia of malignancy: clinical features, diagnosis and treatment]. A hypercalcemia nas malignidades: aspectos clinicos, diagnosticos e terapeuticos. Farias Maria Lucia F de. (Faculdade de Medicina, Universidade Federal do Rio de Janeiro, RJ.. fleuss@hucff.ufrj.br) . Arquivos brasileiros de endocrinologia e metabologia, (2005 Oct) Vol. 49, No. 5, pp. 816-24. Electronic Publication: 2006-01-23. Ref: 64. Journal code: 0403437. ISSN: 0004-2730. Pub. country: Brazil. Language: Portuguese.

AB Hypercalcemia associated with malignancies is reported in up to 20 to 30% of patients with cancer during the course of the disease, and points to a poor prognosis. Symptoms related to the central nervous system, as progressive mental impairment, stupor and coma, predominate. Alterations in kidney function (water-concentrating defect leading to polyuria) and gastrointestinal tract (anorexia, nausea, vomiting) corroborate to dehydration and a further increase in serum calcium. Cancer-induced hypercalcemia may be classified as: 1) local osteolytic hypercalcemia (LOH), due to marked increase in osteoclastic bone resorption in areas surrounding the malignant cells within the marrow space; 2) humoral hypercalcemia of malignancy, caused by the secretion of parathyroid hormone-related protein (PTHrP) by the malignant tumor; 3) ectopic hyperparathyroidism; 4) 1,25(OH)2 D-secreting tumors. Adequate control of hypercalcemia is necessary to give the patient time to respond to anti-cancer therapy. Volume expansion with saline will correct dehydration, improve glomerular filtration and increase urinary calcium excretion, which may be further stimulated by loop diuretics. Intravenous bisphosphonates are the most effective agents to control hypercalcemia, as they block osteoclastic osteolysis and also have antitumoral effects, decreasing bone metastases. New approaches to control the skeletal manifestations of malignancies are anti-PTHrP and anti-RANKL antibodies, osteoprotegerin, and also proteasome inhibitors in the case of multiple myeloma.

=> s l4 and mouth dryness  
L10 0 L4 AND MOUTH DRYNESS

=> s l4 and hyperosmolality  
L11 0 L4 AND HYPEROSMOLALITY

=> s l4 and symptoms  
L12 10 L4 AND SYMPTOMS

=> s l12 and parathyroid hormone

L13 10 L12 AND PARATHYROID HORMONE

=> dup remove l13

PROCESSING COMPLETED FOR L13

L14 4 DUP REMOVE L13 (6 DUPLICATES REMOVED)

=> d l14 1-4 cbib abs

L14 ANSWER 1 OF 4 MEDLINE on STN DUPLICATE 1  
2007004209. PubMed ID: 17200368. Parathyroid hormone

-related protein induces cachectic syndromes without directly modulating the expression of hypothalamic feeding-regulating peptides. Hashimoto Hirofumi; Azuma Yumiko; Kawasaki Makoto; Fujihara Hiroaki; Onuma Etsuro; Yamada-Okabe Hisafumi; Takuwa Yoh; Ogata Etsuro; Ueta Yoichi. (Department of Physiology, School of Medicine, University of Occupational and Environmental Health, Kitakyushu, Japan. ) Clinical cancer research : an official journal of the American Association for Cancer Research, (2007 Jan 1) Vol. 13, No. 1, pp. 292-8. Journal code: 9502500. ISSN: 1078-0432. Pub. country: United States. Language: English.

AB PURPOSE: Parathyroid hormone-related protein (PTHrP) is a causative factor of humoral hypercalcemia of malignancy (HHM) and concurrent anorexia and wasting. Because changes in the expression of hypothalamic feeding-regulating peptides can directly affect appetites and thereby can cause anorexia and wasting, we addressed whether the cachectic syndromes induced by PTHrP rely on the action of hypothalamic feeding-regulating peptides. EXPERIMENTAL DESIGN: Rats were inoculated with a LC-6 human cancer xenograft that secreted PTHrP, and the mRNA levels of the hypothalamic feeding-regulating peptide genes and serum leptin levels were examined before and after the development of HHM by in situ hybridization histochemistry and ELISA, respectively. Some rats were given the anti-PTHrP antibody. RESULTS AND CONCLUSION: The mRNA levels for the orexigenic peptides, such as the agouti-related protein and the neuropeptide Y in the arcuate nucleus (Arc), were significantly increased after the development of HHM, whereas the mRNA levels for the anorexigenic peptides, such as the proopiomelanocortin in the Arc, the cocaine and amphetamine-regulated transcript in the Arc, and the corticotropin-releasing factor in the paraventricular nucleus, were significantly decreased after the development of HHM. Plasma leptin levels were also reduced in cachectic rats, and the administration of anti-PTHrP antibody to the cachectic rats not only improved the cachectic symptoms but also restored the mRNA levels of these orexigenic and anorexigenic peptides, except for orexin. Thus, PTHrP induces HHM and concurrent cachectic syndromes by mechanisms other than directly modulating the leptin or hypothalamic feeding-regulated peptides.

L14 ANSWER 2 OF 4 MEDLINE on STN  
2006057279. PubMed ID: 16444366. [Hypercalcemia of malignancy: clinical features, diagnosis and treatment]. A hipercalcemia nas malignidades: aspectos clinicos, diagnosticos e terapeuticos. Farias Maria Lucia F de. (Faculdade de Medicina, Universidade Federal do Rio de Janeiro, RJ.. fleiuss@hucff.ufrj.br) . Arquivos brasileiros de endocrinologia e metabologia, (2005 Oct) Vol. 49, No. 5, pp. 816-24. Electronic Publication: 2006-01-23. Ref: 64. Journal code: 0403437. ISSN: 0004-2730. Pub. country: Brazil. Language: Portuguese.

AB Hypercalcemia associated with malignancies is reported in up to 20 to 30% of patients with cancer during the course of the disease, and points to a poor prognosis. Symptoms related to the central nervous system, as progressive mental impairment, stupor and coma, predominate. Alterations in kidney function (water-concentrating defect leading to polyuria) and gastrointestinal tract (anorexia, nausea, vomiting) corroborate to dehydration and a further increase in serum calcium.

Cancer-induced hypercalcemia may be classified as: 1) local osteolytic hypercalcemia (LOH), due to marked increase in osteoclastic bone resorption in areas surrounding the malignant cells within the marrow space; 2) humoral hypercalcemia of malignancy, caused by the secretion of parathyroid hormone-related protein (PTHrP) by the malignant tumor; 3) ectopic hyperparathyroidism; 4) 1,25(OH)<sub>2</sub> D-secreting tumors. Adequate control of hypercalcemia is necessary to give the patient time to respond to anti-cancer therapy. Volume expansion with saline will correct dehydration, improve glomerular filtration and increase urinary calcium excretion, which may be further stimulated by loop diuretics. Intravenous bisphosphonates are the most effective agents to control hypercalcemia, as they block osteoclastic osteolysis and also have antitumoral effects, decreasing bone metastases. New approaches to control the skeletal manifestations of malignancies are anti-PTHrP and anti-RANKL antibodies, osteoprotegerin, and also proteasome inhibitors in the case of multiple myeloma.

L14 ANSWER 3 OF 4 MEDLINE on STN DUPLICATE 2  
 2003534792. PubMed ID: 14613038. Treatment of malignancy-associated hypercalcemia and cachexia with humanized anti-parathyroid hormone-related protein antibody. Sato Koh; Onuma Etsuro; Yocum Richard C; Ogata Etsuro. (Department of International Coordination, Chugai Pharmaceutical Co, Ltd, Skizuuoka, Japan. ) Seminars in oncology, (2003 Oct) Vol. 30, No. 5 Suppl 16, pp. 167-73. Ref: 11. Journal code: 0420432. ISSN: 0093-7754. Pub. country: United States. Language: English.

AB Parathyroid hormone-related protein (PTHrP) plays a central role in humoral hypercalcemia of malignancy (HHM), which is one of the most frequent paraneoplastic syndromes. PTHrP produced by the tumor acts through a common PTH/PTHrP receptor to promote bone resorption, inhibit calcium excretion from the kidney, and induce hypercalcemia. Patients with HHM often develop cachexia associated with typical symptoms such as anorexia, malaise, nausea, constipation, polyuria, polydipsia, and confusion. The etiology of the cachexia is not fully understood but is thought to be caused by hypercalcemia and various cytokines such as interleukin-6, tumor necrosis factor-alpha, leukemia inhibitory factor, and others. In this study, we investigated the role of PTHrP in hypercalcemia and cachexia in HHM by using humanized anti-PTHrP antibody. A mouse monoclonal antibody that binds to PTHrP amino acid sequence 1-34 and inhibits PTHrP function has been humanized to create a specific and potent agent for the treatment of patients with HHM. The mouse monoclonal antibody has been shown to have antihypercalcemic activity against nude mice bearing human tumors. Because a mouse antibody is highly immunogenic in human patients, the complementarity-determining regions from the mouse antibody were grafted into a human antibody. The resulting humanized antibody specifically recognizes PTHrP(1-34) and neutralizes PTHrP functions in vitro and in vivo. The humanized anti-PTHrP antibody was administered intravenously to HHM model animals bearing tumors such as LC-6 human lung carcinoma. These animals showed symptoms similar to those of patients with HHM (eg, hypercalcemia and cachexia). The humanized anti-PTHrP antibody-treated animals responded with normalization of blood ionized calcium level through an improvement of bone metabolism and calcium excretion. Moreover, the treated animals also showed an improvement in body weight, ultramotility, metabolic alkalosis, food consumption, water intake, serum phosphorus, and renal function. Consequently, the humanized antibody-treated animals experienced complete resolution of hypercalcemia and cachexia. These results suggest that the humanized antibody would be an effective and beneficial agent for patients with HHM, and that PTHrP is a major pathogenetic factor of hypercalcemia and cachexia in patients with HHM.

L14 ANSWER 4 OF 4 MEDLINE on STN DUPLICATE 3

1995111368. PubMed ID: 7812215. A patient with primary hypoparathyroidism developing hypercalcemia associated with adult T-cell leukemia/lymphoma. Kondo S; Tamura K; Makino S; Yokota T; Ishikawa E; Katakami H; Kohari S. (Department of Internal Medicine, Miyazaki Prefectural Hospital, Japan. ) Leukemia & lymphoma, (1994 Aug) Vol. 14, No. 5-6, pp. 521-5. Journal code: 9007422. ISSN: 1042-8194. Pub. country: Switzerland. Language: English.

AB A 38-year-old woman was admitted to our hospital with symptoms and signs of hypocalcemia in 1977 and a diagnosis of primary hypoparathyroidism was made with a positive Ellsworth Howard test. She was then lost to follow up until 1992 when she returned this time with symptoms and signs of hypercalcemia. An inguinal lymph node was biopsied showing non-Hodgkin's lymphoma, diffuse pleomorphic type and monoclonal integration of proviral human T-cell lymphotropic virus-1 DNA was detected in lymph node cells indicating ATLL. Serum parathyroid hormone-related peptide (PTHrP) was slightly elevated and the tumor cells were positively stained with anti-PTHrP serum. Combination chemotherapy with vincristine, adriamycin, cyclophosphamide and prednisolone was given to the patient with disappearance of the lymphadenopathy and subsequent normalization of PTHrP levels. Interestingly, the signs and symptoms of hypocalcemia reappeared after the treatment requiring replacement therapy with calcium and vitamin D.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	63.22	63.43
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-0.80	-0.80

STN INTERNATIONAL LOGOFF AT 12:58:39 ON 16 DEC 2008